


LECTURE NOTES

For Nursing Students

Pediatric Nursing and Health Care



Teklebrhan Tema
Tsegaye Asres

Jimma University

In collaboration with the Ethiopia Public Health Training Initiative, The Carter Center,
the Ethiopia Ministry of Health, and the Ethiopia Ministry of Education

2003



Funded under USAID Cooperative Agreement No. 663-A-00-00-0358-00.

Produced in collaboration with the Ethiopia Public Health Training Initiative, The Carter Center, the Ethiopia Ministry of Health, and the Ethiopia Ministry of Education.

Important Guidelines for Printing and Photocopying

Limited permission is granted free of charge to print or photocopy all pages of this publication for educational, not-for-profit use by health care workers, students or faculty. All copies must retain all author credits and copyright notices included in the original document. Under no circumstances is it permissible to sell or distribute on a commercial basis, or to claim authorship of, copies of material reproduced from this publication.

©2003 by Teklebrhan Tema and Tsegaye Asres


All rights reserved. Except as expressly provided above, no part of this publication may be reproduced or transmitted in any form or by any means, electronic or mechanical, including photocopying, recording, or by any information storage and retrieval system, without written permission of the author or authors.

This material is intended for educational use only by practicing health care workers or students and faculty in a health care field.

Preface

This lecture note on pediatric and child health is written for nurses at diploma level by considering the epidemiology and the burden of illness on younger age groups. In this lecture note, the authors present concise summaries of the evidence that has been published in the scientific journals and texts which have a similar profile in the Ethiopian context.

Since the authors have had so many years of teaching experience, emphasis is given on those lessons learned from experience.



Acknowledgments

The authors acknowledge The Carter Center (EPHTI) Addis Ababa, Ethiopia for their financial support and encouragement.

Dr. Hailu Nida, assistant professor of pediatric and child health Jimma University, School of Medicine and Sr. Bosna Tebeje, Lecturer, at Jimma University School of Nursing deserve special thanks for their invaluable comments on the lecture note at the intra-institutional work shops in Agaro.

Nega Assefa and Sr. Meselech Asigid both from Alemaya University and Yosief Tsige from Jimma University need to receive warm appreciations for their immense contribution during the inter-institutional work shops in Addis Ababa.

Last but not least, students at Jimma University, particularly in the school of nursing, deserve special thanks for their active participation during the process of testing this manual in the class-room setting.

TABLE OF CONTENTS

Preface	i
Acknowledgement	ii
Table of content	iii
List of tables and figures	viii
Abbreviations	ix





CHAPTER SIX: NORMAL GROWTH AND DEVELOPMENT	76
6.1. Growth	77
6.2. Development	79
Study Questions	84
CHAPTER SEVEN: NUTRITION AND NUTRITIONAL DEFICIENCIES	85
7.1. Normal feeding pattern	86
7.2. Assessment of Nutritional status	89
7.3. Protein Energy Malnutrition	91
7.4. Micronutrient and its deficiency	95
Study Questions	102
CHAPTER EIGHT: ACUTE RESPIRATORY INFECTIONS	103
8.1. Introduction	104
8.2. Management of Children with ARI	105
8.3. Antibiotic Treatment	110
8.4. Home Care	111
8.5. Referral Criteria	112
8.6. Croup or Obstructive Laryngitis	112
8.7 Tonsillitis, Pharyngitis	114
8.8. Chronic Tonsillitis and Adenoditis	115
8.9. Ear Infections	116
Study Questions	118

CHAPTER NINE: CONTROL OF DIARRHEA	119
9.1. Introduction	119
9.2. Types of diarrhea	120
9.3. Why is diarrhea dangerous?	120
9.4. How does diarrhea cause dehydration?	121
9.5. Treating Diarrhea	121
9.6. Prevention of dehydration	122
9.7. Feeding	122
9.8. Oral rehydration therapy	123
9.9. Assessment of a child with diarrhea	124
9.10 Treatment of Diarrhea	129
9.11 To treat diarrhea at home use treatment plan A	130
Study Questions	135
CHAPTER TEN: SYSTEMIC DISEASES	136
10.1. Cardiovascular system (CVS) disorder	136
10.2. Central nervous system Diseases	144
10.3. Epilepsy/ Seizure	147
10.4. Congenital heart Disease	150
10.5. Nephritic Syndrome	153
10.6. Urinary Tract Infection	155
10.7. Tuberculosis	157
10.8. Disease of the Skin	160
10.9. Other problems	163
10.10. Ophthalmic neonatrum	165
10.11. Congenital Syphilis	167

CHAPTER ELEVEN: VACCINE PREVENTABLE DISEASES	168
11.1. Justification for immunization	168
11.2. Immunization schedule	169
11.3. Hazards of immunization	170
Study Questions	174
UNIT TWELVE: EXPANDED PROGRAM ON IMMUNIZATION (EPI)	175
12.1. Strategies of EPI service Delivery	176
12.2. Contraindications	178
12.3. Sterilization	179
12.4. Principles of the immunization program	179
Study Questions	180
UNIT THIRTEEN: COMMON GENETIC PROBLEMS OF CHILDREN	181
13.1. Mental retardation	181
13.2. Cerebral Palsy	183
13.3. Down's Syndrome	184
13.4. Cretinism/ hypothyroidism	186
13.5. Juvenile Diabetes Mellitus	186
Study Questions	188
BIBLIOGRAPHY	189

TABLES AND FIGURES



Abbreviations

ABO	Blood Group Type A,B, and O
AIDS	Acquired Immuno Deficiency syndrome
ALRI	Acute Lower respiratory Infections
APGAR	Appearance, Pulse, Grimace, Activity, and respirations
ARI	Acute Respiratory Infections
BCG	Bacillus Calmette and Guérin
BP	Blood Pressure
CDD	Control of Diarrhea Diseases
CNS	Central Nervous System
CSF	Cerebro Spinal Fluid
CVA	Costa vestibular angle ness
DPT	Diphtheria Pertusis and Tetanus
EPI	Expanded Program on Immunizations
GI	Gastro Intestinal
GUS	Genito Urinary Syndrome
ICP	Intra Cranial Pressure
IM	Intra Muscular
IMCI	Integrated management of child hood illness
IV	Intravenous
NGT	Naso gastric Tube
PaO ₂	Partial pressure of Oxygen
PEM	Protein Energy malnutrition
P'O	By Mouth

PPF	Procaine penicillin Fortified
PRN	When Necessary
RBC	Red Blood cells
RH	Rhesus factor
TB	Tuberculosis
TTC	Tetracycline
VIT A	Vitamin A



and Sweden. In technically advanced countries, the child survival rate is over 97% while in less developed countries over 50% of total deaths are of children under the age of 5, and the average life span is about 35 years. In scientifically advanced countries, only 5% of the total mortality occurs among the under 5; and the average life span is over 70 years. Before the advent of scientific medicine, it was taken for granted that a large proportion of children born alive would die in childhood, and the parents felt it necessary to have many children in the hope that some would survive, that few of them paid much attention to childhood diseases

1.2. Safe mother hood in relation to child health

High Neonatal Mortality Rates are preventable Proven, simple, low cost solutions exist; up to 70% of neonatal deaths could be prevented through intervention for the mother. Impact of mother baby package interventions on neonatal death (WHO mother –baby package 1994).

A. Antenatal Care:

In the first visit

- History taking
- Physical examination

- Grand multiparity
- Preeclampsia, eclampsia
- Cardiac disease
- Renal disease
- Diabetes mellitus
- Multiple pregnancy

B. Labor and delivery Care:

Trained health worker during labor and delivery should attend Mother and baby. Delivery should be conducted under hygienic condition. Clean hand, delivery surface, clean cord-cutting and tying, clean environment, and clean perineum. Skilled attendant at delivery. Recognition of danger signs (mother and baby); treatment and referral as needed.

Age and disease patterns:

C. Essential Newborn Care:

Immediate assessment

Care of the new-born baby should be guided by the following principles

Dry the new-born as soon as possible

Wrap the new-born in a dry towel to prevent heat loss from the body

Place the new-born next to the mother to get the breast and warmth (rooming in).

Cut the cord when it becomes thin and white. Apply

tetracycline eye ointment 1% to both eyes to prevent eye infection. Delay washing of the new-born for 12-24 hours to avoid unnecessary chilling. Nothing should be applied to the cord except gentian violet.

1.3. Age and disease and patterns:

To define the task better, it is important to look in more details at the types of diseases that are prominent at childhood. The following list summarizes the most important diseases that will be encountered at different ages.

- Birth to 1 month-neonatal period
- Obstetric complications and birth injuries
- Asphyxia (failure to breathe at birth)
- Low birth weight babies (prematurity)
- Infection leading to septicemia
- Tetanus (from infected cord)
- Death of mother – the new born baby is likely to die of starvation, (or gastroenteritis, if bottle-fed) unless another woman can be found to breast feed him.

A. First year of life- infancy

- Respiratory diseases- pneumonia, whooping cough, etc.
- Diarrheal diseases (Especially likely if the baby is bottle fed)




before five years of age)It is obvious from this list that most of these child hood diseases are preventable. This why such a great emphasis must be put on preventive programs including nutrition education, immunization and environmental sanitation

1.4. Health Priorities in children

Child health Indicators:

A. Perinatal mortality rate:- the total number of still-births



to the newborn during the first months of life. Immaturity of the infant is the chief cause of these early deaths. Approximately 80 % of infants who die within 48 hours of birth weigh less than 2500 g.

C. Post-natal mortality rate: - The number of deaths over 28 days but under one year of age per 1000 live births.

D. Infant mortality rate: - The number of infant under one year of age dies per 1000 live births. It is the sum of neonatal and postnatal deaths. The primary cause is immaturity and the second leading cause is gastroenteritis, which can be prevented by putting the newborn immediately with the mother and advocating breast-feeding.


E. Child mortality rate: - The number of deaths between 1 and 4 years in a year per 1000 children. This rate reflects the main environmental factors affecting the child health, such as nutrition, sanitation, communicable diseases and accidents around the home. It is a sensitive indicator of socioeconomic development in a community and may be 25 times higher in developing countries compared to developed countries.



B) Major child health programs/activities:

- CDD see the section on IMCI case management of Diarrhea
- ARI see the section on IMCI case management of ARI
- Malaria control see the section on IMCI case management of fever
- HIV control activities (mother to child transmission prevention) see the section in HIV prevention.

C) Antenatal and deliver care:



E) Under Five Clinic:


- The aims and objectives of the under-fives clinic are to provide curative, preventive and promotive health services within the resources available in the country such as:
- Well baby care
- Care in illness
- Adequate nutrition
- Immunization

F) Traditional Practice:

Everyone is greatly influenced by the traditional customs of his family, tribe, and country. Some of these traditional practices are good for health, such as breast feeding or the acceptance of modern medicine, should be supported. Those traditional practices and beliefs, which are bad (cutting of uvula, female genital mutilation etc) need to be gradually changed. This is another important area in which health workers can have a strong influence in improving health.

Study Questions

1. Discuss the components of safe motherhood that can contribute to the reduction of neonatal morbidity and mortality.
2. What are the immediate essential newborn cares?
3. What are the major components of child health program activities?
4. Discuss advantages and disadvantages of traditional practices in relation to child health



CHAPTER TWO

HISTORY TAKING AND PHYSICAL EXAMINATION

Learning Objectives

After studying the material in this unit, the students will be able to:

- Take relevant/proper pediatric history
- Perform systematic physical examination
- Interpret history and physical examination to arrive at differential diagnosis
- Outline Lab. Work up and management

2.1 History taking and approach

A) The Chief complaint: is the symptom the patient presents himself with. It can be stated briefly as e.g. fever, cut wound, or vomiting etc.

B) History of the present illness: this is a chronologic

- Duration of disease onset
- Severity
- Aggravating and alleviating factors
- Associated symptoms
- Any treatment and response to treatment
- History of contact with similar illness
- Relevant pediatric history (like history of immunizations) related to chief complaints or history present illness



well as the current living condition. Poverty and ignorance are major sources of ill health. You may have to educate a poor mother with malnourished baby that the best treatment for her baby is to be breast-fed exclusively till the age of one or more. If a mother feeds the baby food containing unbilled water the baby may get diarrhea. Teach the mother to boil water used for preparing food to infants.



The Symptoms of disease (e.g. respiratory distress, dehydration, pain, anxiety). This will direct your history taking. The healthy looking child whose mother believes he is ill. See how the proposed illness affects the general wellbeing or growth of the child. History taking does not have to be long if he looks well. Inappropriate reaction of the mother the recent history may be irrelevant. Social history may be more important to get.

Abnormal reaction of the child towards
What is happening around him
history may be directed towards CNS

2. Listen to the mother's description of the complaints carefully and get the main symptoms.

- How did the disease start, which symptoms followed?
- Get the time factor. When the disease starts? Has the child had the same kind of symptoms before?
- Ask about the condition of other members of the family?
- How is the child eating at present? How was his appetite in the last month?
- How is he doing between the attacks of illness? Is he weak and inactive or strong and active?
- It may be necessary to obtain the social history in certain conditions such as malnutrition.
- Get the 'story'. Where has the child gone for help before?

What sort of local treatment has he had? What are the mother's beliefs about his illness/disease?

- Always try to evaluate the weight progress of the child. This is best done by looking at the weight chart which the mother preferably should have and bring to outpatients' department at every visit.
- Ask about vaccinations done.
- When the symptoms are vague and nonspecific, e.g. tiredness, abdominal pain etc. review the different systems of the body.

2.2 Review of Systems:

The review of system is essentially the same as in the adult history. It is best organized from the head to down to the extremities. In the child, however, there should be increased emphasis on the symptoms related to the respiratory, gastrointestinal, and genitourinary systems. The high incidence of symptoms and diseases related to these symptoms obligate the interviewer to focus in this area.

To get the important points:

The patient usually comes with his mother and the task is to pick out from all the different information the mother is giving what is important.

- Try to make good contact with her and with the child

- Treat them as human beings who have come for help and advice and do not look upon them as 'cases' only.
- Always believe what the mother tells, but try to be realistic about the importance the symptoms she is mentioning.
- If he is not well nourished and not properly immunized, take the opportunity of a minor illness to prevent major diseases by advice and vaccination.
- Use communication skill (APAC: Ask ,Praise, Advice and check ones understanding)

2.3 Physical Examination:

A) Principles and techniques of physical examination:

The principles and techniques of physical examinations in the case of a small children you should make it habit to undress the child and examine the whole child. To examine the whole body we start with the head and end at feet in older children and adults. In order not to frighten small children it is best to examine things that are uncomfortable or frightening to them last so as not to loose their cooperation. This means the last thing to do in a child is auscultation of the heart, inspection of the ears with an auriscope and inspection of the throat with a throat stick. We use our eyes, ears and hands in addition to a few special items of equipment to perform the physical examination.

1. Chronological steps of physical examination:

This consists of three parts:

a. General appearance:

This is what you observe while examining your patient

The mental state of the patient

- is he acting normally?
- is he confused?
- is he drowsy, stuporous or even comatose? etc.

The general physical state of the patient

- general state of health
- weight and body build
- colors
- respiration
- signs of dehydration
- edema

b. Vital signs:-

These are:

- Temperature
- Pulse rate
- Respiratory rate
- Blood Pressure

The temperature:

All sick children should have their temperature measured (rectally, orally, and axially) The normal temperature is about 37° C. A temperature below 36°C is abnormally low and may be a sign of infection in a small baby.

A temperature above 37.5°C is fever. When there is a fever it usually means an infection is present and you must try to locate the site of the infection and decide whether it needs treatment and with what.

The pulse:-

The pulse can be felt and count in children radically for fifteen seconds multiply by four. In the infant it is sometimes easiest to count the heart rate with the stethoscope apically.

Normal pulse rate:

- babies 100-140 beats per minute
- children 80-100 beats per minute

In fever the pulse rate generally rises. In dehydration the pulse rate may be very rapid and weak.

The respiratory rate:-

Normal respiratory rate:

- < 2 month ,< 60 breath per minute
- 2-12 months < 50 breath per minute
- 12 month-5years< 40 breath per minute

A rapid respiration of 60 or more in a small, feverish child is a very good indicator of pneumonia

c. Anthropometric measurement:

- Weight
- Height/Length
- head circumference
- mid-arm circumference
- Chest circumferences

Weight:

The best way to assess nutritional status is to take body weight. The weight should be charted on a weight chart. Most weight charts have three curves. The upper line shows the average weight of healthy well nourished children and this is an ideal growth curve. The middle shows the lowest weight that is still considered to be within limits of normal and the weights on this line are 80 % of the weights on the upper line. The lower curve shows 60 % of the ideal weight. According to Gomez classification any child whose weight is below this line is marasmic.

Height:

not affected much for the first six months in malnutrition and is therefore more a measurement of longstanding malnutrition.

d. Evaluation of various body system:

For various body system evaluation we use our

- eyes for inspection
- hands for palpation
- ears for auscultation
- HEENT (hair color and texture, pallor, oral lesions, ear discharge, eye discharge, neck swellings...) inspection.
- Chest (Inspection, Palpation/ Percussion, Auscultations)
- CVS (Inspection, Palpation/, Auscultations)
- Abdomen(Inspection, Palpation/ Percussion, Auscultations)
- GUS (Costo-vertebral angle (CVA) tenderness, Suprapubic tenderness, and inspection of external genitalia)
- Musculoskeletal system(joint swelling or tenderness or deformity, bone swelling or tenderness, muscles)
- Integumentary /skin (color, lesions)
- CNS(Level of consciousness, reflexes,(motor and sensory)and meningeal Signs

Study Questions

1. What are the essential components of history taking?
2. What are the principles and techniques of physical examination in a small child?
3. What methods can you use when evaluating the body system?
4. What techniques would you use to get information about the sick child?




CHAPTER THREE



Since the possibility of error is greater in the giving of medication to children than to adults, and since a child's reaction to a dose ordered by a physician is less predictable than adult's reaction, the nurse must be alert to recognize undesired effects of the medication given.

A) Oral administration:

Infants will generally accept the medication put into their







piece of tape to ensure that it reaches the stomach after it is passed. The tip of the catheter



3) Resuscitation:

Respiratory arrest means that there is no apparent respiratory activity. The child will be unresponsive, pale and dead like. Cardiac arrest follows quickly after respiratory arrest as soon as the heart muscle is affected by the anoxia, which occurs. The outcome for the child will depend to great extent on the speed with which resuscitation is began. The steps for resuscitation can be remembered as "A, B, C, D" where A is for airway, B for breathing and C is for circulation and D is for drug administration. These three techniques (clearing the airway, ventilating the lungs, and circulating the blood by cardiac compression) will provide adequate oxygenation to major body organs for an extended period of time (A: clear airway, position, suction), (B: bagging, oxygen administration), (C: Cardiac compression, secure IV line), (D –drugs (epinephrine).

Oxygen administration: Oxygen administration elevates the arterial saturation level by supplying more available oxygen to the respiratory tract. Nursing care must be planned carefully when children are in tents:

- The tent should be open as little as possible so that as high an oxygen concentration as possible can be maintained.
- Oxygen may be delivered to an infant by use of plastic mask. This tight fitting plastic can keep oxygen concentration at nearly 100 %.

- A nasal catheter used with an oxygen flow of 4 L/ min provides a concentration of about 50 %. Most children do not like nasal catheter because it is irritant; assess the nostrils of the infant carefully when using nasal catheter. The pressure of catheter can cause areas of necrosis, particularly on the nasal septum.
- Oxygen must be administered warmed and moistened, no matter what the route of delivery; dry oxygen will dry and thicken, not loosen secretions.
- Oxygen must be administered with the same careful observation and thoughtfulness as any drug.
- If concentrations are too low, oxygen is not therapeutic
- In concentrations greater than those desired, oxygen toxicity can develop.
- If newborns are subjected to a Pao₂ of over 100-mm/Hg for an extended time, retinopathy of prematurely can occur.

4. Administering Enemas:

Enemas are rarely used with children unless a part of preoperative preparation or are required for radiological study.

The usual amount of enema solution used are as follows:

- Infant: less than 250 ml
- Preschools-6(co)5.5(pe)5.nem
-

For an infant:

- Use a small soft catheter (no 10 to 12 French) in place of an enema tip.
- Infants and children up to ages 3 years or 4 years are unable to retain enema solutions, so they must rest on a bed pan or baby pan during the procedure
- Pad the edge of the pan so that it is not cold or sharp.
- Place a pillow under the infants or the young child's upper body for positioning and comfort.
- Lubricate the catheter and insert it only 2 to 3 inches (5cm – 7cm) in children and only 1 inch (2.5 cm) in infants.
- Be certain to hold the solution container no more than 12-15 inch above the bed surface so the solution flows at controlled rate.
- If the child experiences intestinal cramping, clamp the tubing to halt the flow temporarily and wait until the cramping passes before instilling any more fluid
- If the enema solution is to be retained, such as an oil solution, hold the child's buttocks together after administration
- Tap water is not used because it is not isotonic and causes rapid fluid shifts of water in body compartments, leading to water intoxication.
- Normal saline (0.9 sodium chloride) is the usual solution

3.3 Care of child under specific conditions:

Fever: Fever in children is common and can have many causes. In many cases you will make the diagnosis of non-specific virus infection. This may be true, but such a diagnosis is difficult to prove and should never be made without taking a careful history and performing a proper examination in any child with fever. Young children appear to tolerate fever better



General management of children with fever:

- a. If the fever is high (over 39 degree centigrade)
 - Tepid sponging with ordinary water will help to reduce but ice cold water is harmful because it causes constriction of blood vessel in the skin and prevents heat loss.
 - Antipyretics (paracetamol) 10 –15 mg /kg
 - Use a fan to increase water evaporation and hence skin cooling and reduction in body temperature.
- b. Feverish children should never be wrapped in warm blankets. Children must be able to get rid of the heat, otherwise febrile convulsions can be precipitated
- c. children who have fever need to take extra fluids, so make sure that they have plenty to drink.
 - If they are too sick to take, give fluids by NG tube
 - If they are obviously dehydrated, they need intravenous or intraperitoneal infusion
 - Room ventilation
 - Re-assess the child for fever later
 - Attempts should be made to identify and treat underlying causes

B) Coma:

Coma is a state of unconsciousness from which patient cannot be aroused by any stimulation.

- This might be the result of a recent convulsions
- Coma can be caused by local damage to the brain by

3. Turn the patient every 2 hourly
4. Check the bladder regularly for retention
5. Insert NGT for feeding

C) Convulsive seizure: Convulsions are much commoner during the first two or three years of life than at any other period. This is not only due to congenital malformation or perinatal injury to the central nervous system but also the frequency of "febrile" convulsions in response to a rapid rise of temperature at the onset of acute infective illnesses

1. Management of Convulsion in children:

- (A,B,C,D) of life saving measures
- Give 20 % glucose (IV or PO)
- Give Diazepam, (Rectal, IV,NGT) if convulsion does not stop
- Give Phenobarbitone or Phenyntine if convulsion does not stop
- Give General anesthesia

2. Nursing Management during seizure:

- Provide privacy
- Protect head injury by placing pillow under head and neck
- Loosen constrictive clothing's
- Remove any furniture from patient side
- Remove denture if any

- Place padded tongue blade between teeth to prevent tongue bit
- Do not attempt to restrain the patient during attack
- If possible place patient on side

3. Nursing Management after seizure:

- Prevent aspiration by placing on side
- On awaking re-orient the patient to the environment
- Re-assure and calm the patient

3.4 Tracheostomy Care (Care of a child under Tracheostomy):

A tracheotomy is a procedure in which an opening is made into the trachea. When an indwelling tube is inserted into the trachea, the term tracheostomy is used. A tracheostomy may be either temporary or permanent.

A tracheostomy is performed to bypass an upper airway obstruction, to remove tracheobronchial secretions, to prevent aspiration of oral or gastric secretions in the unconscious or paralyzed patient and to replace an endotracheal tube. There are many disease processes and emergency conditions that make a tracheostomy necessary.

Procedure:

The procedure is performed in the operating room or in an intensive care unit, where the patients condition can be well controlled an optimal aseptic technique can be maintained. An opening is made in the second or third tracheal rings. After the trachea is (opened) exposed a tracheostomy tube of appropriate size is inserted.

The tracheostomy tube is held in place by tapes fastened around the patients neck usually, a square of sterile gauze is placed between the tube and the skin to absorb drainage and prevent infection.

Complications:

Early complications immediately after the tracheostomy is performed include:

- bleeding
- pneumothorax
- air embolism
- aspiration
- subcutaneous or mediastinal emphysema
- recurrent laryngeal nerve damage or
- posterior tracheal wall penetration.

Long-term complications include:

air way obstructions due to accumulation of secretions.

-




Tracheostomy tube and skin care:

- Inspect tracheostomy dressing for moisture or drainage
- Wash hands
- Explain procedure to pt.
- Wear clean gloves, remove soiled dressing and discard.
- Prepare sterile supplies (hydrogen peroxide, normal saline, or sterile water, cotton tipped applicators, dressing materials).
- Put on sterile gloves.
- Cleanse wound and tracheostomy tube with sterile applicators moistened with hydrogen peroxide and rinse with sterile saline.
- If old tapes of outer cannula are soiled change it with care and secure with a knot.
- Remove old tapes and discard.
- Use sterile tracheostomy dressing and fit securely to cover the incision.

Study Questions

1. What are the indications for Oxygen administration?
2. What are similarities and difference in giving IM medication to children and adults?
3. Discuss about the indications for NG tube feeding.
4. Discuss about the possible causes of fever in children and its management.
5. Discuss about the care of a child admitted to hospital in state of coma.
6. List the possible causes of convulsion in children.



CHAPTER FOUR

CARE OF THE NEW BORN

Learning Objectives:

After studying the material in this unit, the students will be able to:

- Provide basic delivery room care for the new born
- Identify high risk new born as early as possible and able to undertake basic neonatal resuscitation
- Provide proper nursing care for the normal neonate immediately after birth
- Perform assessment of the newborn before discharge
- Detect and manage the common neonatal problems
- Refer neonates with serious problems as soon as possible to higher facilities

4.1 Care of Normal Newborn:

A) Care at birth:

- Clear air way (suction mouth and pharynx)
- Temperature control (dry and wrap with dry cloth)
- clamping the cord immediately after the birth of baby app. 2.5 cm from abdomen

- provide appropriate eye care to prevent Ophthalmia Neonatorum(TTC eye ointment)
- Identification of newborn with mothers name
- Vitamin K injection to prevent hemorrhagic disease of the newborn
- Measurement (Weight, Height, Head circumference)
- Encourage rooming-in immediately after delivery

Assessment of APGAR score:

- 7-10 well adjusted
- 5-7 mild asphyxia
- 3-4 moderate asphyxia
- < 3 severe asphyxia

B) Care in the Postnatal Ward:

- Handle the baby gentle
- Avoid infection (hand washing)
- Keep the baby warm
- Encourage and initiate exclusive breast feeding -4 months of age
- Encourage care of umbilical stump
- Give immunization (BCG & OPV0)

C) Assessment of the newborn before discharge:

- Take vital sign
- Check for sign infections (eye , cord ,skin)

- Check for birth trauma (cephalohematoma, fracture..)
- Check for anomalies (spina-bifida, imperforate anus, club



Appropriate for gestational age (AGA)- birth weight between and 10th and 90th percentile

Small for gestational age-(SGA) – birth weight below 10th percentile

4.2 Causes of Low Birth Weight (LBW):

Fetal:

1. Fetal distress
2. Multiple pregnancies
3. Erythroblastosis fetalis

Placental:

- i. Premature separation
- ii. Placenta previa

Uterine: Incompetent cervix

Maternal:

1. Pre-eclampsia
2. Chronic illness
3. Infection
4. PROM (premature rupture of membrane)
5. Poly hydrominous >1500ml
6. Nutrition status of the mother

A) Management of low birth weight:

Clean air way

Initiate breathing

Establish circulation

Keep Warm

Administer Vit. K

Administer Silver nitrate or TTC ointment to the eyes

Apply gentian Violet to the cord

Feeding- Breast feeding

NGT feeding

Parental feeding

Treat anticipated problems (infection)

Hypothermia (cold injury). If Rectal temperature is $< 35^{\circ}\text{C}$

Hyperthermia temperature $>38.5^{\circ}\text{C}$

f



Immediate Newborn care:

- Drying, wrapping, close contact with mother
- Immediate breast feeding
- Prophylactic eye care, where appropriate

Routine Newborn Care:

- Exclusive breast feeding, counseling for problems
- Maintenance of body temperature
- Clean cord care
- Early postpartum visit for mother and baby (within 3 days)
- Recognition of danger signs; treatment and referral as needed
- Immunization

Special Newborn Care:

- Special care for the sick children
- Recognition and management of birth asphyxia
- Recognition and management of infection
- Special care for the LBW infant
- Skin – to- skin / kangaroo mother care
- Close surveillance for signs of infections and feeding difficulties



- Anemia
- Drug (narcotic)

Clinical Manifestations:

- Tachypnoea
- Intercostals muscles retraction
- Flaring of Alai nasi
- Cyanosis

Management:

- Resuscitation
- Oxygen administration
- Fluid administration
- Warm - heat

3. Congenital pneumonia

It is caused by aspiration of amniotic fluid or ascending infection.

- Predisposing factors:
- Prolonged Tw(-)-1Tj/aolonxmiBctorsspobed Tachypnoea
-

4.3 Perinatal and infant infections:

1. Neonatal sepsis

Causes:

- Streptococcus
- E. coli
- Premature rupture of membrane (PROM)

Four routes of infection:

- Transplacental
- Ascending infection
- Bacterial colonization
- Environmental

Clinical Manifestation:

- Hypothermia
- Apnea, respiratory distress
- Failure to feed
- Vomiting and jaundice

Management:

Antibiotic (Gentamycin 5-7mg (kg) TID for 14)

Neonatal meningitis: is an infection of brain caused by sepsis or bacteria in the blood stream.

Management:


- Ampicillin 200mg/kg. 24 hr in 4 divided dose
- Gentamicin 5-7 mg/kg/ 24 hrs TID 14 days
- Neonatal pneumonia: is an infection of lung tissue in neonatal period.

Route of infection:

- Transplacental
- Amniotic fluid infection
- Environment
- Instrument

Other Neonatal problems:

-





4.5. Neonatal resuscitation:

During the initial resuscitation efforts, a 100 % oxygen concentration is administered to the neonate. As soon as the infants condition is stabilized, the concentration is adjusted to maintain the PO₂ (partial pressure of oxygen in the blood) within acceptable limits. This adjustment is essential, since elevated pao₂ levels can cause irreparable damage to retinal vessels. Furthermore, high oxygen concentrations can directly injure lung tissue premature infants with immature lungs and eye vessels are at particular risk for two conditions that are a direct result of oxygen toxicity: retrolental fibroplasia and bronchopulmonary dysplasia.

4.6 Fever:

Fever in children is common and can have many causes. In many cases you will make the diagnosis of non- specific virus infection. This may be true, but such a diagnosis is difficult to prove and should never be made without taking a careful history and performing a proper examination in any child with fever.

NB. Remember to ask for any contact with infectious diseases.

Think of malaria and take a blood film. If you still do not have a definite cause for the fever, rule out:

- A. Malaria: one negative blood film report does not exclude malaria
- B. Early measles: look for koplik's spots
- C. Pneumonia: look at the child for flaring of nostrils, rate of breathing, Lower chest in drawing
- D. Otitis media: check eardrums
- E. Meningitis: neck stiffness, irritability
- F. Urinary tract infection: check urine
- G. Tonsillitis: look at the throat
- H. Relapsing fever: take blood film for haemo parasite

4.7 Convulsions:

Convulsions are much common during the first two or three years of life than at any other period. This is not only due to congenital malformation or perinatal injury to the central nervous system but also the frequency of "febrile "convulsions in response to a rapid rise of temperature at the onset of acute infective illnesses

Causes:

1. In the neonatal period the major causes of convulsions are
 - Congenital defect of the brain
 - Cerebral damage occurring during the process of birth from hypoxia or trauma both account for 90 % of the cases.

The remaining 10 % includes:

- infection of the brain (meningitis)
 - hypoglycaemia
 - hyperbillirubinaemia with kernicterus etc
2. during early childhood (until about the age of 6 years)
 - Infections both (intra and extra cranial) are the most frequent causes of convulsions
 - Febrile convulsions between age of 6 months and 3 years and rare after this period
 - At the age of two idiopathic epilepsy begins to accounts for an increased proportion of cases.
 3. at later childhood and adolescence
 - 50 % of recurrent convulsions have idiopathic origin
- The remainders are organic

4.8. Feeding Recommendations during sickness and health:

Up to 4 months of age

- Breast feed as often as the child wants, day and night, at least 8 times in 24 hours.
- Do not give other fluid of fluids
- Expose the child to sunshine for 20 to 30 minutes daily

4 months up to 6 months

- Breast feed as often as the child wants, day and night , at least 8 times in 24 hours
- Add complement foods mashed potatoes softened with milks cereal and legumes mixed with milk
- Give these foods with cup and spoon 1 or 2 times per day in addition to breast feed.
- Expose the child to sun shine for 20 to 30 minutes daily.

6 Months up to 12 months.

- Breast-feed as often as the child wants.
- Give adequate servings of shiro fitfit, Merek fitfit, porridge, made of cereal and legume mixes, mashed potatoes and carot, mashed gommen, egg and fruits.
- Give adequate servings of:Porridge made of cereal and legume mixes. Shiro, kik, merek fitfit, mashed potatoes and carot, gommen,undiluted milk and egg and fruits

- Add some extra butter or oil to child's food
- Give these foods:-3 times per day if breastfed 5 times per day if not breastfed
- Expose child to sunshine

12 months up to 2 years:

- Breast feed as often as the child wants, Give these foods 5 times per day
- Give adequate serving of: porridge made of cereal and legume mixes. Shiro, kik, merek fitifit mashed potatoes and carrot, gommen, undiluted milk and fruits
- Add some extra butter or oil to child food
- Give these foods 5 times per day


2 years and older:

Give family food at least 3 times each day.

Also twice daily, give nutritious food between meals, such as: egg, milk, fruits, kita, dabo

Study Questions

1. List the common causes of low birth weight.
2. Discuss the focus areas to prevent prenatal and neonatal mortality.
3. What are the common causes of infant and childhood mortality?
4. What are the common causes of respiratory distress in newborns?
5. List common routes of infection in neonates?



CHAPTER FIVE

CONGENITAL ABNORMALITIES

Learning Objectives:

- After studying the material in this unit, the students will be able to:
- Define congenital abnormality
- Recognize the common types of congenital abnormalities and check each newborn baby to be sure that there is no congenital abnormalities
- Recognize the recommended treatment for each type of congenital abnormality
- provide appropriate care for children with congenital abnormalities

congenital abnormalities are compatible with intrauterine life, but not with extra-uterine life approximately 15 % of death in the neonatal period care caused by such gross malformations.

5.2 Types of commonest congenital anomalies

1. Cleft lip and palate
2. Club foot
3. Umbilical hernia
4. Pyloric stenosis
5. Phimosi and paraphimosis
6. Spina bifida and hydrocephalus.

5.2.1. Cleft lip and palate

Cleft lip and palate are congenital deformities due to the failure of various parts of the upper lip and palate to fuse in the normal manner. Both may be present together. Cleft lip is operated on about the age of three months, cleft palate about the age of one year before speech defects have developed.

Cleft lip: may be unilateral or bilateral and may be extended up into nostril. The surgeon pares the age and stitches them together.

Preparation: The child should be admitted several days before operation and observed for signs of a cold, as operation should be delayed if this is present and throat swabs should be taken.

The hemoglobin must be 70% before operation is undertaken.



After Care:

Arms must be splinted

Soft feeds are given by spoon placed well back on the tongue and followed by sterile water.(others similar on cleft lip after care)


5.2.2 Club foot

Club foot is a foot which has been fell out of shape or position in uteri and can not be moved to uncorrected position.

The two most common types of club foot are:

Talipes equinovarus

Talipes calcaneovalgal



Treatment and Responsibility of the Nurse

Treatment should be started as soon as possible. Delay



5.3 Umbilical Hernia:

An umbilical hernia is a protrusion at a portion of the intestine through the umbilical ring, muscle, and fascia surrounding the umbilical cord due to imperfect closure or weakness of the umbilical ring. The clinical manifestation is a swelling at the umbilicus which is covered with skin. This protrudes when the infant cries or strains. It can be reduced by gentle pressure over the fibrous ring at the umbilicus. The contents of the hernia are small intestine and omentum.

Treatment and Responsibilities of the Nurse

- Most small umbilical hernias disappear without treatment, but large ones may require operation. These hernias rarely cause incarceration or strangulation of the bowel.
- If the physician orders the hernia to be taped, the hernia is first reduced by gentle pushing the abdominal contents back through the umbilical ring.
- A 2 inch strip of adhesive tape can be applied to the abdomen over the skin covering the umbilical hernia.
- Usually operation is not done on an umbilical hernia unless it becomes strangulated, enlarges or persists to school age.
- Postoperative nursing care requires no special technique. The child may be as active as his desire. A normal diet and fluid may be given; pressure dressing applied at the time of operation must be kept clean and dry to prevent wound contamination.

5.4. Congenital Hypertrophic pyloric stenosis

This is a common surgical condition of the intestinal tract in infancy. It occurs most frequently



Prognosis: The prognosis is excellent complete relief follows successful surgical repair. The mortality rate is low, proceeded operation is undertaken before the infant has become too dehydrated and malnourished.

5.5. Phimosis and paraphimosis

Phimosis is a tight for skin, i.e, a fore skin or prepuce which has a very small opening, so that it may interfere with the passage of urine and can not be drawn back over the glans penis for cleansing. In severe case this will prevent the scape of urine, so that the fore skin is distended during mictaration, and the infant screams with pain and strains during the act.

Paraphimosis, a condition due to a tight foreskin being pushed up and constrict the circulation, causing swelling of the parts below so that the foreskin cannot be pushed down again, and requiring operative treatment.

Treatment:

- Circumcision, cutting away the foreskin
- After this operation the wound must be kept clean, antiseptic dressing applied, and the parts cleansed after passing of the urine.
- Mild cases of phimosis can be relieved by gradually stretching the foreskin by pulling it back over the glans penis, so that operation is avoided.

5.6 Spina Bifida

Spina bifida is a malformation of the spina in which the posterior portion of the laminae of the vertebrae fails to close. It may occur in almost in any area of the spina but most common in lumbosacral region.

It is the most common developmental defect of the central nervous system occurring in about one of 1000 newborn infant.

The three types are:

1. spinabifida occulta in which the spinalcord and meninges are normal, the defect being only of the vertebrae,
2. Meningocele, in which the meninges protrude through the opening in the spinal canal.
3. Meningomylocele, in which both the spinal cord and the meninges protrude through the defect in the bony rings of the spinal canal. Meningomylocele is the most serious one.

Spina bifida occulta:

The majority of patients with spina bifida occulta have no symptoms. Some may have a dimple in the skin or growth of hair over the malformed vertebra. There is no need for treatment unless neurological symptoms indicate that the

defect is greater than was thought. If there is possibility that the spinal cord may be involved in the defect, surgical treatment is indicated.

Meningocele:

On examination the newborn infant is found to have a defect on the spinal column large enough to protrude through the opening. There is generally no evidence of weakness of the legs, the infant straitens and kicks in normal manner, or if lack of spincter control, though this is difficult to a certain in the newborn. The prognosis is excellent if surgical correction is done on this defects. Hydrocephalus may be an associated finding or may be aggravated after operation for a meningocele.

Meningomyelocele:

In this condition an imperfectly developed segment of the spinal cord as well as the meninges, protrudes through the spina bifida in the lumbosacral region. There may be a minimal weakness to a complete flaccid paralysis of the legs and absence of sensation in the feet.

Operation removes a cosmetically unacceptable deformity, prevents infection and in many instance improves the neurological deficit since obstruction is removed from the nerve path ways.




The obstruction inflow cerebrospinal fluid may be due to one of several causes.

Congenital mal development of the ventricular foramina.

Neoplasm may be present or fibrous residue making it may occlude the reabsorptive surface.

Hemorrhage from trauma may cause hydrocephalus in the young infant.

The accumulation of fluid in the ventricles generally enlarges the infants skull, since the suture airhcles generally enlarges



If pressure areas develop, great care must be taken to prevent infection.

When lifting head must be supported in order to prevent trauma.

Postoperative care:

- The temperature, pulse and respiration should be observed every 15 minutes until the infant is reactive.
- Signs of increased intracranial pressure (irritability, bulging of the fontanel, lethargy, vomiting, elevated systolic blood pressure, change in the pulse and respiration rate, change in body temperature must be recorded and reported.
- If temperature is elevated tepid sponging may be used or aspirin may be given.
- IV fluid must be given slowly until the infant can be fed orally.
- Mucus from nose and mouth should be aspirated to prevent the danger of aspiration.
- The child should be turned at least every two hourly.

5.8 Neonatal Jaundice:

Jaundice is a yellow discoloration of the skin, sclera and mucous membranes caused by high serum bilirubin. Bilirubin is the main pigment formed in humans during the catabolism

of the hem component of hemoglobin. Either excessive production or defective elimination of bilirubin causes jaundice.

Physiologic jaundice resolves spontaneously by the end of first week in term infants and 2nd week in premature infants.

Serum bilirubin is usually under 12 mg/dl in term infants and less than 15mg/dl in premature babies.

Jaundice within 24 hrs of birth; conjugated bilirubin over 12mg/dl; or jaundice beyond 2nd week of life is pathological

Causes:


- ABO/RH incompatibility
- Red cell membrane defects
- RBC enzyme defects
- Polycythemia
- Infection

Management of pathological jaundice

- Enhancing conjugation by induction of glucuronyl transferase with phenobarbiton
- Changing insoluble bilirubin to a water soluble form by phototherapy
- Removing excess bilirubin by exchange transfusion.

Study Questions

1. Discuss the role and responsibility of nurse in identifying and caring for a child with congenital abnormality at birth
2. List the most common congenital abnormalities that can be detected at birth
3. What measures would you take concerning parents who had a baby with congenital abnormality?



CHAPTER SIX

NORMAL GROWTH AND DEVELOPMENT

Learning Objectives

After studying the material in this unit, the student will be able to:-

- Identify the difference between growth and development
- Recognize stages of normal growth and development
- Detect deviation from normal growth and development measures to be taken
- Use growth-monitoring chart to assess nutritional status of under 5 children
- Recognize needs of the growing child

Definition: Growth means increase in size, Development means increase of function. Growth and development go together but at different speeds.

The fetus: during the first trimester the main body systems are prepared. This is mainly the stage of development, when the body systems become more efficient



The figures below are average figures for normal growth.

Table 1. Normal growth and development

Age	Weight	Length	Arm circumference
Birth	3.5 kg	50 cm	
½ year	7 kg		
1 year	10 kg	75 cm	12.5cm
2 year	12 kg		
3 year	14 kg		
4 year	16 kg	100 cm	
5 year	18 kg		

Head circumference:



the rest of the year another 3cm. If the head circumference at birth is 34 cm it will be 46 cm at the age of 12 months. The head circumference is measured by taking the greatest distance around the forehead and the back of the head above the ears (maximal fronto-occipital circumference)

Arm circumference: The circumference of the middle of the upper arm remains nearly constant from 1-5 years.

Tooth eruption: Children start teething at the age of 6 months. A new teeth appears approximately every month. This makes the number of teeth roughly equal to the age in months minus six. At the age of 6 years the permanent teeth start to appear.

6.2 Development:

actors promoting development:

Nutrition: Good nutrition is the base for normal growth and development. Unlike most other organs in the body, the brain is not fully developed at birth. The first six months of life are extremely important as the brain may suffer for the rest of life, if the child is not getting enough food. A malnourished child is often tired and apathetic and not interested in learning new things that will promote normal development.

Emotional Support: The first years of life are the most important for the development of the child. It is very important to realize that a child is a growing and developing human being and he ought to be treated very carefully with love and respect by everyone so he can develop in harmony. He needs full emotional support. Here are six important aspects of what every child needs.

Love: a child who does not feel loved will not develop properly, and will not learn as quickly as normal children. Instead he becomes sad, lonely and no longer interested in what goes on around him.

Security: a child needs to feel safe. He can only feel safe if his parents show that they love him and take good care of him. He must know that his parents will look after him and help him, that they will feed him when he is hungry, play with him and keep him happy and comfortable. The love and security the child gets from his parents and family helps him to feel friendly to people outside his family when he grows up.


Acceptance as individual: the young child needs to know that his mother and family love him for what he is. They should not compare him with other children and tell him that he is slow to do this or that, or that he is not as good as some other child. They should show him that they respect him as an individual with his own likes and dislikes, that they realize that he is different, as all children are different.

Recognition of achievement: the young child needs to know that his parents are happy and pleased when he has learned to do some thing new. His parents help him to do things and encourage him in his achievement. Wise and consistent use of authority: children need to know what they can and what they cannot do. Parents must teach children how they are expected to behave.

Independence: as the child grows he needs to be allowed to decide more and more things alone.

Playing: another important factor contributant fa6(J-1s3wr72rtS81s3wr72c[8 m Tw[:

inkthin(g(o1.



- What he can do in the way of moving around (motor development)
-



Table 2. Summary of normal developmental milestones:

Average age	motor development	Language & social behaviors
1 month	can lift head when prone	can fix with eyes, often smiles
3-6 months	good head control	can follow an object with eyes, & play with hands
6-9 months	can sit unsupported	grasp actively, makes loud noise
9-12 months	able to stand	understand few words, tries to use
12-18 months	able to walk	grasp small objects with thumb and fingers
2 years	able to run around	can say several words, or even as much as he wants some sentences
3 years	actively plying, is clever in climbing and jumping	starts talking much



CHAPTER SEVEN

NUTRITION AND NUTRITIONAL DEFICIENCIES

Learning Objectives:

After studying the material in this chapter the student will be able to:

- Identify the common causes of infant and child morbidity and mortality
- State the most common complication of malnutrition and their management
- List the most common forms of malnutrition in the developing Countries and their management
- State the purpose of screening and identify children who need appropriate management
- Perform weighing of babies under 3 years of age and interpret the weight according to weight for age chart
- Classify nutritional status of children and manage accordingly including nutrition education.

7.1 Normal feedings patterns:

The main rule for feedings children

Only breast milk during the first 4 months start adding porridge at 4 months.

Add some protein to every meal after 4-5 months of age.

Plain porridge is not enough for more than a few weeks.

Add fruits and vegetable to at least one meal every day start when the child is 4 months old.

Give a child at least three meals /day. Infants may require four to six small ones.

Continue breast feeding until the child is able to manage on other foods probably this will be at 18-24 months of age.

Use the common local carbohydrate food and add a suitable protein food.

Prepare balanced meals. Make sure that the added protein is well mixed with the staple

Carbohydrate food and that both are given in a suitable amount.

A) Breast-Feeding:

The food for babies is undoubtedly human breast milk. Nurses and midwives should encourage and support mothers to breast feed even if it is only achieved for a few weeks. However, if a mother has decided, for what ever the reason,

to use a milk formula method of bottle feeding she should be given the same support, understanding, and help as the mother who breast feeds.

Good Reasons for Breast-feeding:

1. Promote bonding between mother and child.
2. Many infections are far less common in breast fed babies.
3. The fat in human milk is better absorbed than the fat in cow's milk
4. The danger of giving an over concentrated or a too diluted formula milk is avoided
5. The risk of eczema, asthma, and other allergic disorders is considerably reduced
6. Low birth weight and pre-term infants thrive best on breast milk
7. It helps for mental development

and of the need of extra protein and calories while breast feeding.

4. Practical demonstration and discussion with a breast-feeding mother

Management of Breast-Feeding:

Immediately after birth, while the mother still in the delivery room, all mothers should have the opportunity to hold the baby for 10-15 minutes quietly and contently. If the mother tends to breast-feed, she should suckle her infant for 1-2 minutes at each breast. Skin-to-skin contact and early suckling are important at this time in promoting "bonding" and to encourage the release of the prolactin and oxytocin hormones which stimulate milk secretion and help the uterus to contract.

B) Weaning Food:

The age at which solid foods can be introduced to the infants depend upon individual family circumstances. Most authorities agree, however, that it should be some time between 3 and 6 months of age for the normal healthy infant.

Principles of weaning:

1. Start with one food at one feed time, offering a small quantity only
2. Introduce new foods one at a time

3. Introduce second meal after 3-4 weeks. Include iron containing foods, e.g. liver, green vegetables
4. As solid foods increased and milk volumes reduced, remember to offer dilute fruit juice or water from a cup to infant atleast twice a day particularly during hot weather.

7.2 Assessment of Nutritional Status:

Growth Monitoring:

The first step in appraising the growth of a child is by comparing the child with others of the same age and sex. A child, generally, is healthy if she/he grows well and gains weight. When a child is not growing well, he is probably not healthy. The causes for this ill health could be infection or inadequate food intake. Because of these, the growth of a child will slow down or stop months before clear signs of protein energy malnutrition (PEM). Thus, measuring the growth of a child helps to understand if the child is healthy or not. Growth monitoring is particularly important for follow-ups for children under 3 years of age. You are advised to refer growth-monitoring chart.

How to interpret the growth chart:

The first sign of protein energy malnutrition (PEM) is growth failure. Weighing a child regularly on a growth chart and understanding the direction of the growth line are the most important steps in detection of early malnutrition.

Table 3. Gomeze classification of Nutrition Status

Weight for age/reference weight	Edema present	Edema absent
60-80%	Kwashiorkor	Underweight
< 60%	Marasmic Kwashiorkor	Marasmus

It is very important to follow subsequent measurements and plotting, to watch the direction of the line showing the child's growth.

Table 4 Interpretation and the findings of growth charts

Indication	Child's condition Good gaining weight	Danger sign Stagnant	Very Dangerous Losing weight
Indication of the growth Monitoring chart			
Interpretation	Child is growing well	Not gaining weight Find out why Poor nutrition infection	Losing weight may be ill, needs care
Intervention	Complement the mother	Instruct the mother, support her	Careful counseling return soon, admit or refer

7.3 Protein Energy Malnutrition

The most common form of malnutrition in the world today is that associated with diarrhea, as well as an inadequate diet in the young children. The side effects of malnutrition include hypoglycemia, hypothermia, hypotonic and mental apathy. Severe malnutrition (Kwashiorkor and Marasmus) contributes to high mortality and morbidity of children less than five years in developing countries and to mental and physical impairment in those who survive.


A) Kwashiorkor:

The most acute form of malnutrition is generally found in a child of 10-14 months who has had an excessive carbohydrate diet containing relatively little protein. Edema and enlargement of liver may later mask a slight loss of weight. Coldness of the extremities is well marked and the child is miserable but apathetic when anorexia and/or diarrhea set in, there is loss of weight in spite of edema. This marasmic stage may be due to malabsorption rather than deficiency in caloric intake. Chronic cases show depigmentation of skin and hair, with the hair losing its luster, becoming straight, dry and sparse.

Marasmus:

This condition, seen in children whose weight is markedly

below normal for their length is described as state of starvation. A general deficiency of protein and energy has occurred, leading to severe wasting of subcutaneous fat and muscle tissue. It often occurs between the ages of 6 months and 18 months. The marasmic child appears as a wizened old man in appearance, with loss of most fatty tissue, shriveled buttocks and emaciated limbs. Many of the signs of kwashiorkor, such as edema, sk



- If he is apathy prevent hypoglycemia by IV glucose administration
- Treat skin or chest infection with an appropriate antibiotic

Later treatment

- Give him high protein foods by mouth later or as soon as he eats
- Give him iron mixture after the first week of treatment and continue until his hemoglobin is normal
- Explain the mother the importance of diet treatment and make sure that she understands that food not medicine is curing him
- Caring for a malnourished child:
- In addition to diagnosis of PEM there are five things we have to know
- How severely malnourished is he?
- His weight chart is useful
- What other diseases has he?

Many children with PEM have infections and some lack Vitamins

Which of the six rules of good nutrition have been broken?

- Breast feeding until at least 18 months
- Start supplementary diet at the fourth month
- Add protein foods to supplementary diet
- Give children four good meals a day

- Give protective foods to children over four months old (fruits, vegetables)
- Sick children need more feeding

Why have the rules been broken?

- Poverty?
- Lack of knowledge?
How could his mother feed him?

Prevention of malnutrition

The following are some of the important approaches in prevention of malnutrition in children.

- Encouraging and protecting breast-feeding
- Improving weaning practices by using local foods.
- Screening of infants and children at risk for malnutrition and give them
- Special attention
- Providing nutrition rehabilitation especially in households.
- Integrate nutrition education into primary health care (PHC).
- Ensure regular supervision in clinics and visiting in homes by inspection and growth monitoring

E) WHO/IMCI Classification of PEM/Anemia

Look and feel;

- Visible severe wasting
- Look for palmer pallor (some/severe)

- Look for edema on both feet
- Determine weight for age

Table 5. Classification of malnutrition:

Signs	classification
Severe visible wasting or Severe palmer pallor or Edema on both feet	severe malnutrition/severe anemia
Some palmer pallor or Very low wt. For age	anemia or very low weight

7.4 Micro-nutrient/Vitamins deficiencies:

Vitamins are groups of organic substances not synthesized by the human body but essential as catalysts of cellular metabolism and thus for normal growth. Since the body does not manufacture vitamins small amounts must be included in the diet. Some are soluble in fat and are ingested in dietary fat (vitamin A, D, E and K), and some are water soluble (Vitamin B complex and Vitamin C).

Vitamins; Functions and Deficiency

1. Vitamin A

Normal growth, normal vision, normal reproduction
 Maintenance of epithelial cell structure and function
 Immunity to infection

Deficiencies results in:

- xerophthalmia, (night blindness, conjunctiva dryness, Bitot spots, Keratomalacia, and even eyeball perforation and blindness)
- Increased risk of infections (Viral is more).

Excess results in:

- Raised intracranial pressure, irritability, dry skin, hair loss, brittle bones

2. Thiamin (B1)

Carbohydrate metabolism, decarboxylation

Synthesis of fatty acid and Ribose (for RNA)

Deficiency results in: Beriberi manifested by

- Anorexia, emotional disturbances. Parasthesia, weakness, gastrointestinal symptoms
- Dry beriberi (peripheral neuropathy, mental confusion, nystagmus)
- wet beriberi (biventricular cardiac enlargement, systemic venous hypertension, bounding pulse)
- Infantile beriberi (acute cardiac failure)

3. Riboflavine (B2)

- Coenzyme in oxidative –reduction reaction

Deficiency results in;

- Angular stomatitis, cracking and fissuring of lips
- Glossitis, papillary atrophy
- Scrotal or vulvae dermatitis
- Photophobia, corneal vascularisation
- Anaemia, hair loss , ataxia
- Personality changes, retarded intellectual development

4. Niacin (nicotinic Acid)

- Oxidative –reduction reactions , fat synthesis, glycolysis
- Deficiency results in: Pellagra (dermatitis, diarrhea, dementia)
- loss of weight, poor appetite, sore mouth, indigestion
- insomnia, confusion
- skin erythema, pruritus, discoloration, flaking

5. Pyridoxine (B6)

- Coenzyme in amino acid metabolism and
- Muscle glucogen phosphorylase

Deficiency results in:

Infants: hyperirritability, convulsions, weakness anemia, and dermatitis

6. Folic Acid

- Co-enzyme in pyrimidine and purine synthesis (for DNA,RNA)



9. Vitamin E (tocopherol):

- Antioxidant (protects against free radicals)
- Preserve cell membrane integrity

Deficiency result in:

- Hemolytic anemia, skin changes
- Encephalomalacia

10. Vitamin K

Synthesis of coagulation factors

Deficiency results in:

- Coagulopathy: haematuria, hematomas, and hemorrhagic disease of newborn
- Hemolytic anemia may be caused by the water soluble form of vitamin K

Iron deficiency Anemia:

Anemia refers to a deficit of red blood cells or hemoglobin in the blood. It is the most frequent hematological disorder encounter in children.

Etiology:

- 1. Blood loss
- 2. Impairment of red blood cell production

a. Nutritional deficiency

- Iron deficiency
- Folic acid deficiency

Vitamin B12 deficiency

b) Decreased erythrocyte production:

- Pure red cell anemia
- Secondary hemolytic anemia's associated with infection, renal disease, and chronic disorders Aplastic anemias

Invasion of bone marrow by

A, Leukemia

B, Tumors

3. Increased erythrocyte destruction

- a. Drugs and chemicals
- b. Infections
- c. Antibody reaction
- d. Burns
- e. Poisons including lead poisoning
- f. Abnormalities of the red cell marrow
- g. Hemolytic disease of the newborn
- h. Abnormal hemoglobin synthesis
- e.g. sickle cell disease

Clinical Manifestations:


1. The condition may be acute or chronic
2. Early symptoms
 - a. Listlessness
 - b. Fatigability
 - c. Anorexia

3. Late symptoms
 - a. Pallor
 - b. weakness
 - c. tachycardia
 - d. Palpitation
4. Eventual symptoms



Study Questions

1. How can a nurse contribute to decrease morbidity and mortality among under five children?
2. What are the important points to observe during screening in under five clinics?
3. What are some of the important approaches in preventing malnutrition among under five children's?
4. During growth monitoring9(enti444(n2sewTh)-0.6(ve chbh(ve chat(im(ve coh)-0.6ortant)T



CHAPTER EIGHT

ACUTE RESPIRATORY INFECTION (ARI)

Learning Objectives:

After studying the material in this chapter the student will be able to:-

- Describe the magnitude of acute respiratory infection morbidity and mortality among under five children in Ethiopia
- Identify and use simple clinical signs to classify and treat a child with acute respiratory infection (ARI)
- Distinguish the clinical manifestations of ARI among children under 2 months of age and older children
- Identify danger signs of ARI in infants and children and take appropriate action
- Identify and use commonly available and effective antibiotics against Common etiologic agents of ARI.
- Give advice to the mother on home treatment of ARI.



drawing develops. The presence of lower chest in drawing means that the child has severe pneumonia and is at a higher risk of death. Because of the high probability of bacterial pneumonia and the reliability of these two diagnostic criteria, there is a strong justification for the empirical use of antibacterial based on simple diagnostic criteria.

8.2 Management of Children with ARI

The management of children with ARI comprises three essential steps.

- Identify children who should be examined for possible pneumonia
- Identify the case of pneumonia (case classification)
- Institute the appropriate treatment

Examine children <5 for possible pneumonia if they have cough or difficult breathing

Classification and Diagnostic Criteria:

It is necessary to distinguish between infants under 2 month of age and older children because the etiology and clinical manifestations of pneumonia are different in these age groups.

a) Children 2 months – 4 years old

In these age groups, children with cough or difficult breathing are classified in to 4 groups.



higher chest compliance in young infants, the presence of severe chest in drawing is required to classify an infant below 2 months as having severe pneumonia. There is no pneumonia if there is no danger signs, no severe chest in drawing or not fast breathing. It should be noted that young infants could become sick and die very quickly from pneumonia. For this reason, any young infant who has signs of pneumonia (chest in drawing or fast breathing) is classified as having a severe pneumonia.

c) Treatment of acute lower respiratory disease (ALRI)

Children below 5 years of age with ALRI are divided into 3 therapeutic categories. Children with very severe disease or severe pneumonia must be referred to a health center or a hospital with inpatient capacity where adequate treatment can be given/provided. Before referral, a first dose of antibiotic should be given and fever treated if present.

Children with pneumonia must be treated for 5 days on ambulatory basis with a first line antibiotic recommended by the program. Other problems (fever, ear infection etc) should also be treated if present. The child should return after two days for reassessment. If the child shows sign of a very severe disease or severe pneumonia she/he must be referred. If the child does not show signs of a very severe disease or severe pneumonia but did not improve, different antibiotics

should be given for other two days. If the child still does not show improvement, s/he should be referred.

c. A child having cough for more than 30 days (chronic cough) should be referred for further assessment.

Children with no pneumonia must not be treated with antibiotic. Advice to the mother should be given on home care and other problems assessed and treated.

Table 6 a. ARI Clinical Chart: children < 2 months

Signs, cough plus	Classification	Treatment
Stopping feeding well or Convulsion Abnormally sleepy or difficult to walk or Strider in calm child or Wheezing or Fever or low temperature or Severe chest in drawing or fast breathing or	Serious bacterial infection	- Refer urgently to hospital - Give first dose of antibiotic - Refer urgently to hospital - Give first dose of antibiotic -Treat fever if present
Fast breathing but no chest in drawing	Pneumonia	-Advise mother to give home care - Give an antibiotic - Keep young infant warm


Table 6b. ARI Clinical Chart: children 2 months – 4 years of age

Signs	Classification	Treatment
<ul style="list-style-type: none"> - Not able to drink or - Convulsion or - Abnormally sleepy or difficulty to awake or - Stridor in calm child or - Chest in drawing 	Severe pneumonia or very severe disease	<ul style="list-style-type: none"> - Refer urgently to hospital/ admission - Give first dose of antibiotic - Treat fever if present - If cereberal malaria is possible give ant malarial drug(s) - Refer urgently to hospital - Give first dose of antibiotic - Treat fever if present
<ul style="list-style-type: none"> - Fast breathing 	Pneumonia	<ul style="list-style-type: none"> - Advise mother to give home care - Give an antibiotic - Treat fever if present
<ul style="list-style-type: none"> - Not fast breathing and no chest in drawing 	No pneumonia	<ul style="list-style-type: none"> - If there is cough for more than 30 days refer for assessment - Advise mother to give home care - Treat fever if present

- amoxicillin
- ampicillin
- cotrimoxazole and
- procaine penicillin

8.4 Home Care

Mothers of children treated for pneumonia should return home



Mothers of young infants should also be told to keep the infants warm and clear the baby's nose if it interferes with feeding.

8.5 Referral Criteria:

Children with a very severe disease or severe pneumonia must be referred to HC or hospital. Refer only if it is expected that the patient will receive better care. If this is not the case, the child should be treated with the available antibiotics and needless referral avoided.

Explain to the mother the reason why the child should be referred. If you think the mother will not take the child to hospital or the referral will be delayed, whatever the reason, you should take the following steps:

- If timely referral is likely give first dose of antibiotic
- If there is long referral time give additional doses
- If referral is uncertain give full course

Clinical Features:

- A) The illness usually follows a cold or accompanies measles. Then the child suddenly develops a croupy cough and inspiratory stridor as a result of obstruction in the area.
- B) In some children a large red and infected epiglottis can be seen.
- C) In some the condition becomes worse, retraction of intercostal space, the supraclavicular space and even of the ribs themselves become more evident. The child become restless, pale, shows obvious sign of air hunger

Treatment:

- A) Steam: liquefies the dry secretions and results in marked improvement. A bed sheet over an infant bed makes a perfect steam tent
- B) A humidified oxygen if available can be life saving
- C) Make sure the child drinks enough
- D) Prednisolone 2mg /kg/day decreases edema
- E) Antibiotics: PPF 50,000u/kg with sulphdimidine orally 100mg/kg/day in 3 divided doses for 5 to 7 days
- F) Trachetomy or intubation can be lifesaving
- G) Mild sedation with phenergan 1mg/kg/ day can be of help in:

8.7 Tonsillitis, Pharyngitis:



Complications:

- A. Otitis media
- B. Peritonsillar abscess (in older children only)
- C. Cervical lymphadenitis may develop into abcess
- D. Rheumatic fever or acute glomerulonephritis after



3. General symptoms of chronic infection (tiredness, poor appetite) are common

Indication for Adenoidectomy or /and tonsillectomy (over 3 years only)

- a. Recurrent otitis media, tonsillitis, cervical adenitis= several times a year
- b. Peritonsillar abscess
- c. Enlarged adenoid should be removed
 - i. if giving rise to recurrent infection
 - ii. if there is loss of hearing
 - iii. if nose breathing is continuously observed

8.9 Ear Infections:

Ear infections in children are classified into 3 categories:

- Mastoiditis
- Acute ear infection
- Chronic ear infection


A child with painful or tender swelling behind the ear is classified as having Mastoiditis. In a young infant the swelling might be above the ear.

- The child should be urgently referred after a first dose of antibiotic



Study Questions

1. What steps would you take for a child with a very severe disease or severe pneumonia before referral?
2. What advice would you give to the mother for home treatment of ARI before she leaves the health unit?
3. If you think the mother will not take the child who needs referral or if the referral will be delayed, what steps will you take?
4. What are the criteria for referring a child with ARI?



CHAPTER NINE

CONTROL OF DIARRHEA DISEASE

Learning Objectives:

After studying the material in this chapter the student will be able to:-

- Identify different forms of diarrhea and their management
- Recognize different causes of diarrhea and its prevention
- Assess and classify a child with diarrhea for dehydration and treat as indicated
- Teach parents and community on the ways of preventing diarrhea among children
- Recognize different forms of a solution to prevent and treat dehydration

9.1 Introduction:

Diarrhea means passing of three or more loose or watery stools in a day. Diarrheal disease is among the leading causes of morbidity and mortality among children < 5 years of age in Ethiopia. Diarrhea is most common in children, especially those between 6 months and 2 years of age. It is also common in babies under 6 months who are not breast-fed.

9.2 Types of diarrhea:

Diarrhea can be classified by its duration as acute or persistent and dysentery

- Acute diarrhea – starts suddenly and may continue for several days but < 2 weeks
- Persistent diarrhea lasts >2 weeks and may vary from day to day
- If the stool contains blood the diarrhea is called dysentery

9.3 Why is diarrhea dangerous?

Acute diarrhea causes death because of dehydration

Dysentery causes death because of a number of severe and potentially fatal complications occurring during dysentery such as

Intestinal perforation

- Toxic mega colon
- Convulsions
- Septicemia
- Prolonged hyponatremia
- Diarrhea is worse in person with malnutrition. Diarrhea can also cause malnutrition and make it worse because
- Nutrients are lost from the body during diarrhea
- Nutrients are used to repair damaged tissue rather than for growth

- A person with diarrhea may not be hungry
- Mothers may not feed children during diarrhea or even for some days after diarrhea stops

To prevent malnutrition, food should be given to children with diarrhea as soon as, they eat it.

9.4. How does diarrhea cause dehydration?

The body normally takes in water and salts it needs through drinks and food. When the bowel is healthy, water and salts pass from bowel into the blood. When there is diarrhea, the bowel does not work normally. Less water and salts pass into the blood, and more passes from the blood into the bowel. Thus, more than the normal amount of water and salts passed in the stool results in dehydration. Dehydration also can be caused by a lot of vomiting, which often accompanies diarrhea.

9.5. Treating Diarrhea:

The most important measures in treating diarrhea are to:

- Prevent dehydration from occurring if possible
- Treat dehydration quickly and well if it does occur
- Feed the child

9.6. Prevention of dehydration:

- Drinking extra fluids as soon as diarrhea starts
- Home fluids as well as ORS if available
- Food based fluids containing salt are most effective. eg. Rice water
- If fluids do not contain salt, give food that is salted to taste
- If dehydration occurs, the child should be treated with ORS at health institution
- In severe dehydration, first treat with IV fluid, but ORS should be used in addition as soon as the patient can drink
- ORS alone when the signs of severe dehydration are gone

9.7 Feeding:

Feeding during diarrhea episode provides nutrients that the children need to be strong and grow

- Breast-fed children should be given the breast more frequently
- Other children should be given their usual feedings
- Children of 6 months or older should be given small amounts of nutritious and easily digestible foods more frequently
- After the diarrhea has stopped, an extra meal each day for 2 weeks to help the child regain weight lost during the illness


9.8 Oral rehydration therapy:

Recommended home fluids

- Breast milk
- Cow milk
- Cereal-based gruel (eg. 'Muk', 'attimit')
- Soup ('shorba')



- Pour the water into the container. Mix well with a clean spoon until the powder is dissolved
- Taste the solution so that you would know its taste like salt
- Then give the child frequent small sips out of a cup or spoon. If the child vomits, wait for 5-10 minutes, then continue giving ORS slowly





When going through the assessment chart:

Ask the following questions:

- For how long has the child had diarrhea?
- How many liquid stool, has the child passed in 24 hrs?
- Is there any blood in the stool?

Has the child been vomiting?

- If so, has the child vomited more than a small amount?
- How often has the child vomited in 24 hrs?

Is the child able to drink?

If so, is the child thirsty than usual?

Look for the following conditions

- What is the child's general condition?
- Is the child:
 - Well and alert?
 - Unwell, sleepy or irritable?
 - Very sleepy, floppy or unconscious?
 - Having fits?

Does the child have tears when he or she cries?

Are the child's eyes normal, sunken or very dry sunken?

Are the child's mouth and tongue wet, dry or very dry?

Is the child's breathing normal, faster than normal, very fast and deep?

When water is offered to drink, is it taken normally, eagerly or is the patient unable to drink?

Feeling for the following conditions:

When the skin is pinched:

- Does it go back quickly?
- Does it go back slowly?
- Does it go back very slowly? (>2 seconds)
- Pinch the skin over the abdomen and thigh
- Can the pulse be felt? (Normal 100-120/min) faster, or weak?
- Is the fontanel normal? Depressed or very depressed in young infants.
- Ask for other problems than dehydration:
- Ask about blood in the stool

If blood is present, treat for 5 days with an oral antibiotic recommended for shigella in your area.

- See the child again after 2 days if:
 - Under 1 year of age
 - Initially dehydrated
 - There is still blood in stool
 - Not getting better

If the stool is bloody after 2 days, change to a second oral antibiotic recommended for shigella in your area. Give it for 5 days.

If diarrhea has lasted at least for 14 days:

- Refer to hospital if:
- The child is under 6 months old
- Dehydration is present (refer after treatment of dehydration)

Otherwise, teach the mother to feed her child as in plan A

Look for severe under nutrition If the child has severe under nutrition:

- Do not attempt rehydration: refer to hospital for management
- Provide the mother with ORS solution and show her how to give it
- A teaspoon every 1– 2 minutes for less than 2 years child
- Frequent sips from a cup for an older child




9.10. Treatment of Diarrhea

Decide on appropriate treatment:

After the examination, decide how to treat the child

- if the child has any of the signs in the column labeled “for other problems” specific treatment is needed in addition to treatment given for dehydration
- if there is blood in the stool and diarrhea for less than 14





Rule 3: Take your child to the health worker if the child is not getting better

The mother should take the child to a health worker if the child does not get better in 3 days or develops one of the following:

- Passes many stools
- Does not eat or drink normally
- Has fever
- Is very thirsty
- Vomits repeatedly
- There is blood or mucus in the stool
- Is not passing urine for > 6-8 hours
- Seems not getting better

Children should be given ORS solution at home if:

They have been on treatment plan B or C

They cannot return to the health worker if the diarrhea gets worse

Show the mother how much ORS to give after each loose stool and give her enough packets for 2 days.

Table 9. Amount of estimated ORS solution.

Age	Amount of ORS to give after each loose stool	Amount of ORS to provide for use at home
< 24 months	50-100ml	500 ml/day
2-10 years	100-200 ml	1000 ml/day
10 years or more	As much as wanted	20000/day

Describe and show the amount to be given after each stool using a local measure

- Give a teaspoonful 1-2 minutes for a child < 2 years
- Give frequent sips from a cup for an older child
- If the child vomits, wait for 10 minutes, then give the solution more slowly (a spoonful ORS every 2-3 minutes)

Use the patient's age only when you do not know the weight

- If the child wants more ORS than shown, give more
- Encourage the mother to continue breast-feeding
- For infants < 6 months children who are not breast fed, give 100-200 ml clean water during this period

After 4 hours, reassess the child using the assessment chart, then select plan A,B or C to continue treatment

- If there are not signs of dehydration, shift to plan A
- If signs indicating some dehydration are still present repeat plan B, but start to offer food, milk and juice as described in plan A
- If signs indicating severe dehydration have appeared, shift to plan C

If the mother must leave before completing plan B

- Show her how much ORS to give to finish the 4 hours treatment at home
- Give her enough ORS Packets to complete rehydration,

- and for 2 more days as shown in plan A
- Show her how to prepare ORS solution
 - Explain to her the three rules in plan A:
 1. To give ORS or other fluid until diarrhea stops
 2. To feed the child
 3. To bring the child back to the health worker if necessary

Table 10. Treatment Plan-C To Treat Severe Dehydration Quickly

Age	First give 30 ml/kg IV	Then give 70 ml/kg
Infants (<12 months)	1 hour	5 hours
Older	30 minutes	2 ½ hours

Guidelines for rehydration therapy for severe dehydrations

Start IV immediately

If the patient can drink, give ORS by mouth while the drip is set up

Give 100 ml/kg Ringers lactate or normal saline in divided dose as above.

Repeat once if radial pulse is still very weak or not detectable

Reassess the patient every 1-2 hours. If hydration is not improving give the IV more rapidly.

Also give ORS about 5 ml/kg/hr as soon as the patient can drink; usually after 3-4 hours (infant) or 1-2 hour (older pts)

Reassess after 6 hours (infant) or 3 hours (older patients) and choose the appropriate plan (A,B,C)

If there are no signs of dehydration use treatment plan A

If some of the signs of dehydration are still present but the child is improving, give ORS for another 6 hours specified in the treatment PlanB.

If the signs of dehydration are worse or remain unchanged rehydration therapy with treatment plan C should be continued.

In particular attention should be given to:

The number and volume of stool


The extent of vomiting

The presence and changes in the sign of dehydration

Whether the rehydration fluid (oral and or IV) is being successfully given and is adequate amount.

Study Questions

1. What are most important points to be considered in treating diarrhea?
2. How does diarrhea cause dehydration?
3. What important measures should be taken to prevent dehydration in children with diarrhea?
4. What steps would you follow when assessing a child with diarrhea?
5. What important pieces of advice would you give to the mother for home treatment of diarrhea?
6. How would you treat a child with severe dehydration?



CHAPTER TEN

SYSTEMIC DISEASES

Learning Objectives

After studying the material in this unit the student will be able to:-

- Identify different forms of systemic diseases affecting children
- Manage different forms of systemic diseases
- Prevent complications arising from systemic diseases
- Evaluate the effectiveness of care and treatment
- Identify danger signs of systemic diseases and take appropriate action

10.1. Cardiovascular system (CVS) disorder

Most heart diseases in young children are congenital. Older children are more likely to have acquired heart diseases such as rheumatic fever, endomyocardial fibrosis. The heart may also be affected in many systemic disorders i.e. infections, malnutrition, anemia etc.

Symptoms and signs related to cardiovascular diseases:

Cyanosis is a bluish discoloration of the skin as result of reduced amount of hemoglobin in the circulating blood. Cyanosis can best be detected under the fingernails or on the mucus membranes of the mouth (lips, under side of the tongue).

Clubbing of fingers and toes often occurs. One of the main causes of this is chronic under saturation of the blood with oxygen.

Signs of Cardiac Failure:

- Tachycardia-rapid pulse
- Tachypnea-rapid respiration
- Dyspnea-shortness of breath
- Edema and other signs of raised venous pressure
- Fatigue and failure to thrive
- Arrhythmia-irregular heart beat
- Systolic and more frequently diastolic murmurs
- Cough
- Orthopnea

Management:

Any child with congestive heart failure should be referred to hospital whenever possible. In all cases where you have to start treatment:

- check weight of the child ,record the pulse and respiration carefully at 2 hours intervals and indicate the exact time of any drugs given.

Treatment:

- A. bed rest
- B. sedation (phenobarbitone 5mg/kg/day in 2-3 divided dose im or rectally)
- C. Oxygen if available
- D. Diuretics 10 mg in stat in acute cardiac failure.
- E. Digitalization is most important


In order to achieve effective blood levels quickly a digitizing dose is calculated and given over 24 hours.

1/2 of the calculated dose is given initially

1/4 8 hours later

1/4 after another 8 hours

Then continue with the maintenance dose, which is usually $\frac{1}{4}$



Pathophysiology:

Anatomically the following principal defects may be found.

- a. Abnormal communication in the heart or between big vessels

- Atrial septal defect

- Ventricular septal defect

- Patent ductus arteriosus

In these due to the highest pressure in the left heart, there is a shunt from left to right heart with an increased blood load in lesser circulation. In this group of disease there is no cyanosis.

- b. Congenital obstruction of the blood flow

- pulmonary stenosis

- aortic stenosis (coarctation of the aorta)

- c. Combination of abnormal communication and stenosis

- (pallot's disease is one example)

Clinical Features:

Besides the above-mentioned symptoms, failure of normal growth and development, repeated attacks of respiratory tract infections, and a loud murmur is usually present.

Management:

- A. Any child with congestive heart failure should be referred to hospital whenever possible. In all cases where you have to start treatment:
 - check weight of the child ,record the pulse and respiration carefully at 2 hours intervals and indicate the exact time of any drugs given.
- B. Respiratory tract infections are readily treated with penicillin in normal doses.
- C. Give chloroquine in case of fever after taking blood film.
- D. Treat anemia if present.
- E. Arrange a well-timed comprehensive vaccination program.
- F. Give prophylaxis against subacute bacterial endocarditis

Prognosis: Many children with congenital heart disease die in

Pathogenesis:

Rheumatic fever usually follows about 2 weeks after an infection of the throat or skin with beta-hemolytic streptococci. This is due to a specific reaction of tissues, mainly the heart and the joints, to the streptococcal toxins.

Clinical Features:

Painful swelling of one or more big joints (knee, ankle, elbow, shoulder) may last for one day or longer, subside and another joint may then be affected (rheumatic polyarthritis)

Fever

malaise

rheumatic carditis (heart become enlarged murmur develops and sign

Of congestive heart failure may occur.

1) Relapses are very likely to occur

Treatment/ management:

A. bed rest particularly in pa

Congestive Heart Failure:

Congestive heart failure occurs when the cardiac output is inadequate to meet the metabolic need of the body and results in accumulation of excessive blood volume in the



Pathophysiology:

- a) For any number of reasons, cardiac output is inadequate to meet the oxygenation and nutritional requirement of vital organs
- b) various compensatory mechanisms occur (Tachycardia)
- c) Cardiac output decreases further as a compensatory mechanism fail
- d) There is diminished blood return to the heart, with venous congestion and a rise in venous pressure.

Diagnostic evaluation:

a. Palpation (may have weak peripheral pulse)

Auscultation (gallop rhythm, cardiac murmur may or may not be present)

Chest x-ray (cardiomegally may be present)

Nursing Care:

1. Administer digoxin as ordered to improve myocardial efficiency
2. Provide uninterrupted rest to reduce energy requirement.
3. Administer diuretics as prescribed to remove accumulated sodium and fluid and restrict sodium intake.
4. Administer oxygen therapy to improve tissue oxygenation.
5. Provide foods to meet calorie requirement of the child
6. Practice careful hand washing technique to decrease the dangers of infection

7. Monitor vital signs frequently and report any significant changes to observe signs of disease progress or response to treatment

10.2. Central nervous system Diseases

Meningitis: Meningitis is an inflammation of the meninges (membranes surrounding the brain and spinal cord) and is caused by a viral, bacterial or fungal organisms. Meningitis is further classified as aseptic, septic and tuberculosis meningitis. Aseptic meningitis refers either viral or other causes of meningeal irritation such as brain abscess or blood in the subarachnoid spaces. Septic meningitis refers to meningitis caused by bacterial organisms such as meningococcus, Staphylococcus, or influenza bacillus. Tubercles meningitis is caused by the tubercle bacillus.

Meningeal infections generally originate in one of two ways either through the blood stream as a consequence of other

2. See that the patient's bowel and bladder are emptied
3. Position the patient on one side with back towards the physician
4. The thighs and legs are flexed as much as possible to increase the space between the spines of the vertebrae for easier entry into the subarachnoid space
5. Small pillow is placed under the patient's head to maintain the spine in horizontal position
6. Assist the patient to maintain the position to avoid sudden movement, which can produce trauma
7. Instruct the patient to breathe normally, because hyperventilation may lower an elevated pressure

Post procedure Care:

1. Instruct the patient to lie prone 2-3 hours to reduce the leakage of CSF.
2. Encourage increased fluid intake to reduce the risk of post procedure headache.
3. Check the puncture site for any leakage/bleeding

Examination of the CSF:

CSF should be clear and colorless. Pink blood-tinged or grossly bloody CSF may indicate a cerebral contusion, laceration, or subarachnoid hemorrhage. Sometimes with a difficult lumbar puncture the CSF initially is bloody because of local trauma but then becomes clearer.

Usually specimens are obtained for cell count, culture, glucose and protein. The specimen should be sent to the laboratory immediately because changes will take place and alter the result if the specimens are allowed to stand.

10.3 Epilepsy/ Seizure

Epilepsy/ Seizure is the recurrent occurrence of convulsion or fit.

Causes:

1. Idiopathic:

- Genetic defects
- Developmental defects

2. Acquired:

- Brain hypoxia
- Fever (childhood)
- Head injury
- CNS infection

Clinical Manifestations:

- A, Few seconds before convulsion begins the patient has a warning sign (aura) such as
- Strange pain in the abdomen or
 - Strange movement of the arm or

His may be affected Immediately after warning sign
patient become completely unconscious:
Falls to ground suddenly or
Falls into open fire
Muscles are tightly clenched

This is the first part of fit or convulsion and lasts only a few seconds (Need formatting)

Following this

All muscles contract and relax quickly
The tongue may be bitten
Urine and feces may be passed

This is the second part of fit or convulsion and the convulsion lasts only few minutes.

These jerky movements are called Convulsions and are diagnostic of major or Grand mal epilepsy.

When convulsions are prolonged or repeated the condition is known as "Status epilepticus". Such convulsions are very exhausting and unless controlled may lead to patient death.

During the third part of the fit or convulsion

Patient lies quietly
Muscles are relaxed
He is still unconscious

This part lasts from a few minutes to an hour or more. Then

Patient return to consciousness

He may have bad headache

Remembers nothing of the fit

Often feels very sleepy

B, Some patients do not show the first or second part of the fit, and suddenly become unconscious for only a few seconds. The eye may stare but see nothing and these are the 'lesser fit of epilepsy (petit Mal)".

Medical Management:

1. Phenobarbitone from 30 mg daily to 120 mg three times a day or
2. Phenytoin sodium from 45 mg daily to 180 mg three times a day may be given

Nursing Management during seizure:

Provide privacy

Protect head injury by placing pillow under head and neck

Loosen constrictive clothing's

Remove any furniture from patient side

Remove denture if any

Place padded tongue blade teethes to prevent tongue bit

Do not attempt to restrain the patient during attack

If possible place patient on side

Nursing Management after seizure:

Prevent aspiration by placing on side

Administer medication as ordered to control the seizure

Remove hard toys from the bed to protect the child from injury during convulsion

Do not give anything by mouth during convulsion

Place the child where he can be watched closely to observe for recurrent seizures

On awaking re-orient the patient to the environment. Re-assure and calm the patient

10.4 Diseases of genitourinary tract

- Acute Glomerulonephritis:

Glomerulonephritis refers to inflammation of the kidneys caused by an antigen antibody reaction following an infection in some part of the body. Acute glomerulonephritis is predominantly a disease of childhood and is the most common type of nephritis in children.

Etiology:

1. Antigen antibody reaction secondary to an infection elsewhere in the body.
2. Initial infection of (upper respiratory system or skin) most frequently a beta hemolytic streptococcus and other bacteria's and viruses.

Pathophysiology:

1. Antibodies produced to fight the invading organism also react against the glomerular tissue
2. The antigen antibody combination results in an inflammatory reaction in the kidney.
3. General vascular disturbances, including loss of capillary integrity and spasm of arterioles, are secondary to kidney changes and are responsible for much of the symptomatology of the disease.

Clinical Manifestations:

1. Onset is usually 1-3 weeks after the onset of the initiating infection
2. Sign and symptoms
 - Decreased urine output
 - Blood or brown color urine
 - Edema result in most patients
 - Hypertension present in 50 % of patients (in 4-5 days of illness)
 - Variable fever and mild headache
 - GI disturbances (anorexia and vomiting often with loin pain).

Diagnostic evaluation:

1. Urinalysis (decreased output, hematuria, high specific gravity, proteinuria, white cells, casts) may be reported

2. Blood urea nitrogen and creatinine often elevated
3. Sedimentation rate –elevated
4. Chest x-ray may show pulmonary congestion and cardiac enlargement

Complications:

1. Hypertensive-encephalopathy (restlessness, stupor, convulsions, vomiting, severe headache, visual



10.5 Nephritic Syndrome:

Nephrotic syndrome refers to a common complex characterized by edema, marked proteinuria hypercholesterolaemia, and hypoalbuminaemia. Although there are many types of the diseases lipid nephrosis is the most common in children.

Etiology: The exact cause is not known, but the symptom complex results from large loss of protein in the urine, too great to the body to replenish by albumin synthesis.

Pathophysiology:

1. Protein, especially albumine, leaks through the membrane and is lost in the urine
2. Plasma protein decreases as protein uria increases
3. Over production of aldestrone causes retention of sodium and results in accumulation of fluid in the inertial spaces
4. Decreased gamaglobine results in increased susceptibility for infection
5. Generalized edema is responsible for (respiratory distress, GI symptoms, loss of body tissue, malnutrition.)

Clinical Manifestations:

1. Insidious onset of edema
2. GI symptoms (vomiting, anorexia,)
3. Severe recurrent infection

4. Marked edema
5. Profound weight gain
6. Decreased urine out put during the edematous phase
7. Nephrotic crisis (albuminur



10.6 Urinary Tract Infection:

Urinary tract infection refer to an infection within urinary system. The lower urinary tract (urtetha,bladder,or the lower portion of the ureters) or the upper urinary tract (upper portion of the ureters, or kidney) or both may be involved.

Etiologic agents:

1. GI flora Organisms responsible for 75 % of all cases.
 - E.Coli (most common causative agent)
 - Proteus
 - Aerobacter
 - Entrobacter
 - kelbsella
 - Psudomonus
2. Streptococci and staphylococci causes most other cases.
3. Contributing causes
 - obstruction
 - infection elsewhere in the body
 - poor perineal hygiene
 - Short female urethra
 - Catheteriztion

Path physiology:

Inflammatory changes occur in the affected portion of the urinary tract

Inflammation results in urine retention and stasis of urine in the bladder

There are inflammatory changes in the renal pelvis and throughout the kidney when this organ is involved:

The kidney may become large and swollen

Eventually, the kidney become small, tissue is destroyed, and renal function fails.

Incidence:

1. Most common renal disease in children
2. More common in females than in males, ratio 6:1 except in the neonatal period, when both sexes are equally affected.

Clinical Manifestations:


1. Fever may be accompanied by chills or convulsion
2. Anorexia and general malaise
3. Urinary frequency, urgency, dysuria
4. Dull or sharp pain in the kidney area
5. Vomiting
6. Failure to thrive in infancy

Diagnostic Evaluation:

1. Urinalysis
2. Urine culture
3. Renal concentrating ability decreases
4. Urological and radiological studies

Nursing Care:

1. Obtain mid-stream urine specimen whenever possible and send to Lab. Immediately for examination
2. Administer antibiotics as prescribed by the pediatrician



Most children infected with *M. tuberculosis* are asymptomatic and are only identified a positive tuberculin skin reaction. Tuberculin skin testing consists of the intradermal injection of 5 tuberculin units of purified protein derivatives

Definition of a positive Mantoux skin test in children:

A reaction greater than or equal to 5mm is POSITIVE in:

- Children in close contact with person who have known or Suspected infectious tuberculosis
- Children suspected to have tuberculosis disease including those
 - With clinical evidence or a consistent chest radiography or
- Children who are immuno suppressed or who have HIV infection

A reaction greater than or equal to 10 mm is POSITIVE in:

- Children younger than 4 years of age
- Children at risk of disseminated tuberculosis including those with chronic diseases like malnutrition, diabetes mellitus, chronic renal failure
- Children born in areas where tuberculosis highly prevalent
- Children exposed to adults at risk of tuberculosis, including adults who are HIV infected, or live in poor living condition etc.

A reaction greater than or equal to 15 mm is POSITIVE in:

Children over 4 years of age with no risk factors .Previous

immunization with BCG does not alter these definitions. The presence of clinical manifestations distinguishes tuberculosis disease from tubercle infection. A chest radiography usually service to distinguish children with the disease from those with the infection. Because the sputum of children with pulmonary tuberculosis is usually negative for mycobacterium, either by acid-fast stain or culture, these children are non infectious.

Extrapulmonary tuberculosis: including cervical adenitis, tuberculosis meningitis, and miliary tuberculosis occurs in app. One quarter of cases of children with tuberculosis disease. Infants and young children who immunocompromised or malnourished are at risk of the serious and often fatal form of miliary or tuberculosis meningitis. Children with tuberculosis infection (without evidence of disease) should receive isoniazid prophylaxis for 9 months. Children with pulmonary tuberculosis are usually treated with a 6-month regimen consisting of isoniazid, rifampin, and pyrazinamide for the first 2 months and isoniazid and rifampin the remaining 4 months. In areas where isoniazid and rifampin resistance is prevalent, a fourth drug should be added to the regimen (usually ethambutol, or streptomycin).

10.8 Disease of the Skin

Burns: Burns are a frequent form of childhood injury. A second degree burn of 10 % or more of the body surface in a child younger than 5 year or a second degree burn of 15 % or more of the body surface in a child over 1 year is considered a very serious injury. The effects of burn are not limited to the burn area.

Causes of Burn in Children:

1. Burns from hot water
2. Burns from open fire
3. Electrical burns
4. Acid or alkali burns of mouth and esophagus
5. Chemical burns of the skin
6. Burns inflicted upon the child as a result of child abuse

Clinical Manifestations:

The characteristics of burn wounds are classified as follows

A . First degree burns involves superficial epidermis; the skin

painless to touch. Symptoms of shock (rapid pulse, subnormal temperature, pallor, prostration, low BP) appear soon after the burn.

2. Symptoms of toxemia (Prostration, fever, rapid pulse, cyanosis, vomiting, edema) may develop within 1-2 days after initial burn. These symptoms may progress to coma or death.

3. Burns of the upper respiratory tract result in symptoms of upper respiratory tract obstruction resulting from edema and inflammation of the glottis, vocal cords, and upper trachea.

Treatment:

The objective of treatment are to:

- replace fluid loss from burn surface
- maintain circulation
- prevent renal failure
- prevent or treat infection
- aim toward early repair of the burn wound
- restore the child to the best possible state of physical and psychological functioning

Complications:

Acute:

- Infection
- wound sepsis
- pneumonia

- Generalized lymphadenopathy, hepatosplenomegaly
- Oral candidiasis, recurrent diarrhea, otitis media.
- Neurological deficits-encephalopathy
- Cardiac involvement with myocardial dysfunction.

Prevention of HIV Infection:

The public should be accurately and well informed


Avoiding sexual transmission of HIV

Use of HIV free Blood or blood products

IV drug users should not share syringes and needles

Disinfecting of the equipment appropriately or blood products

sfu(ssi)6.8(on 3 0 IV)TJ/TT5 0 7.98 .12 640 Tm0 Tc40 Tc f



Complications:

The common complication is corneal ulceration with resulting opacity and partial or complete loss of vision. The extent of the handicap depends on the duration and severity of untreated condition.

Treatment:

The treatments of Ophthalmia neonatorum include: prompt antibiotic injections and antibiotic eye ointments or drops to prevent eye damage and isolation of the infant.

Responsibility of the nurse:

The must be sure that:

1. The drops are instilled within the lids during the treatment
2. Prevent the infant from rubbing the infected eye
3. The purulent discharge must irrigate and removed frequently by directing the flow of fluid from the inner cantus outward.
4. Extreme care must be taken that no drops of the return flow splash into the nurse's eyes.
5. Antibiotic eye ointments are administered to all newborns immediately after birth as prophylaxis.

10.11 Congenital Syphilis

Congenital syphilis is a Transplacental infection of the fetus before the fourth month of pregnancy by the spirochete *Treponema pallidum*. It may result in miscarriage, still-birth, or manifest itself in early infancy.

Clinical features:

- Rhinitis with purulent or even bloody discharge
- Lesions on the (large blisters on palms and soles) and mucocutaneous junctions (mouth, nostrils anogenital)
- Signs of systemic involvement: hepatosplenomegaly, prolonged jaundice, and anemia.
- Painful swelling of long bones with characteristic x-ray changes
- Positive serological test for syphilis in mother and child

Management

1. PPF 100,000u/kg/day im for 10 days
2. Check the whole family for syphilis

Prevention:

All measures which decrease the incidence of acquired syphilis in adults

serological tests for syphilis in all pregnant women is desirable.

CHAPTER ELEVEN

VACCINE PREVENTABLE DISEASES

Learning Objectives:

After studying the material in this unit the students will be able to:-

- Identify vaccine preventable childhood diseases
- Recognize different forms of antigens used to protect against vaccine preventable diseases
- Recognize contraindications and adverse effects of immunization
- Immunize children at appropriate age
- Teach parents and care givers to immunize their children against vaccine preventable diseases

11.1 Justifications for immunization:

Of every 1000 children born in the world, about:

- grow up crippled by poliomyelitis
- die of neonatal tetanus

- 20 die of whooping cough
- 30 or more die of measles and its complication

Yet effective vaccinations against the disease exist.

The purpose of immunization is to protect against infectious diseases before they attack any individual. Immunization is the cost-effective method of infectious disease control. Once an immunization program has been established, it must continue or the disease will return to affect large numbers of unprotected individuals. The aim is to immunize in the first year of life against those infections that cause severe disease in infants and children; and to follow up it up with reinforcing (booster) inoculations and additional vaccines according to age. The most serious diseases of infants are whooping cough, (Pertussis), diphtheria, tetanus, tuberculosis, measles, poliomyelitis.

11.2. Immunization Schedules:

Immunization schedules are determined by:

The immunogenicity of the vaccine: good immunogenic vaccine (measles, BCG) need only one injection to produce long lasting immunity, weak immunogens (DPT, cholera)

The epidemiology of the disease-when the disease occurs at an early age, immunization must be given early tuberculosis (BCG), polio vaccine, pertusis).

11.3. Hazards of Immunization:

a. Complications of immunization:

- Normal toxicity or reactivity- fever, malaise, local swelling
- Bacterial contamination-septicemia, abscess, tuberculosis
- Allergic reaction-anaphylaxis

b. Contraindications to immunizations:

1. Inactivated vaccine

- Major febrile illness
- Major intercurrent infections
- Severe reaction to previous dose
- Specific contraindication to pertusis and DPT

2. Attenuated (live) vaccines

- Major febrile illness
- Major intercurrent infections
- BCG to tuberculin individuals-severe local react


3. BCG Vaccination:0.05 ml of vaccine is injected

intra dermal in the newborn

- 0.1 ml is used in all other ages
- Recommended site- just above the insertion of the deltoid muscle
- If tuberculin positive individuals are given BCG
- the skin reaction will be more rapid-noticeable in 3 days
- the inflammatory reaction will be more intense and likely to ulcerate

Complications of BCG vaccinations:

- Local abscess formation and ulceration
- Regional lymphadenitis
- Local granulomatous lesions
- Hypertrophied scars and keloids



Indication for measles vaccination:

- All children in measles endemic area
-



Study Questions

1. What are the common vaccine preventable diseases?
2. What are the adverse effects of vaccinations and measures to be taken?
3. List the most common contraindications for vaccination?
4. Discuss the importance of vaccines.
5. List the types of vaccines commonly used in developing countries?



CHAPTER TWELVE

EXPANDED PROGRAM ON IMMUNIZATION (EPI)

Learning Objectives:

After studying the material in this chapter, the student will be able to:

- Recognize the six vaccine preventable diseases and their annual incidence rate in Ethiopia.
- State the objectives of EPI and its target groups
- Identify vaccine schedule for each target diseases and provide vaccinations as scheduled.
- Realize the dangers of unsterile vaccination equipment and use appropriate sterilization methods before using it.

Communicable diseases are among the major causes of child mortality in Ethiopia. The estimated annual incidence rates per 100,000 live births for six EPI target diseases are as follows:

- Measles 400
- Pertusis 400



Table 11. Child immunization scheme

Contact	Age of child	Vaccine
1.	At birth	BCG and OPV Zero
2.	6 Weeks	DPT1 and OPV1
3.	10 Weeks	DPT2 and OPV2
4.	14 Weeks	DPT3 and OPV3
5.	9 Months	Measles

The immunization schedules for the six "EPI" antigens are given in the tables above. These tables reflect the ideal situation for immunization. Often mothers bring their children at less frequent intervals than those suggested here. If the child is first seen at a later stage than indicated in the immunization scheme, the immunization is started with vaccines 1,2,3 and 4, given with four weeks interval. If the child is 9 months or older, the 5th injection, measles is given four weeks after DPT3. The vaccination scheme should be started on all children under 2. BCG and measles should be given to all under 5. BCG and measles may then be given at the same time. Tetanus Vaccination: some women have certificate that they had DPTs. They need only one injection at each of their two first pregnancies.

Table 12. Type of Immunization, dosage and route for administration.

Vaccine	Route of administration	Dose
BCG	Intradermal injection	0.05 ml 0.1 ml after 1 year of age
Polio (OPV)	By mouth	2 drops each time
DPT	Intramuscularly injection into the thigh	0.5 ml each time
Measles	Subcutaneous injection left upper arm	0.5 ml

12.2. Contraindications:

There are only three contraindications to immunizations.


1. Give no BCG vaccine to a child with clinical AIDS
2. Do not give DPT2 and DPT3 to a child who has had convulsions or shock within three days of the previous dose
3. Do not vaccinate in the presence of severe illness. Vaccinate at later occasion when the child has recovered. Children with diarrhea should be given OPV. This dose should not be counted.

12.3. Sterilization:



Study Questions

1. What are the strategies used to achieve the objectives of EPI in Ethiopia?
2. What are the dangers of using contaminated syringes and needles during vaccination and what are the steps to be used to prevent it?
3. The vaccine preventable diseases estimated to account for one third of infant and child disability and mortality in Ethiopia. What factors do you think contributed to this?



CHAPTER THIRTEEN

COMMON GENETIC PROBLEMS OF CHILDREN

Learning Objectives:

After studying the material in this unit, the student will be able to:-

- Identify the most common genetic problems of children
- Manage children with different genetic problems
- Support parents of children with genetic problems
- Help children with mental retardation to be trained basic skills

13.1. Mental Retardation:

If a child is significantly retarded in his psychomotor development (milestones)

we suspect subnormal intelligence and speak of mental retardation. Several degree of mental retardation is seen. Severe, moderate, mild. In very mild case there may be only minimal brain damage giving rise to slight delay only and less than optimal intelligence may only become obvious in school.





Clinical Features:

1. Spastic paralysis if the lesion unilateral or spastic paraplegia if the lesion bilateral
2. Mental retardation may or may not be present
3. Ataxia is prominent
4. Deafness and blindness can occur as well
5. Squint is common

Management:

1. Regular exercise under the guidance of physiotherapist help in preventing deformity and contracture
2. Simple home training must be initiated
3. Parents of such children support and reassurance

Prevention:

- Proper antenatal and perinatal care,
- early recognition of meningitis etc

13.3. Down's Syndrome:

Down's syndrome is a chromosomal abnormality involving an extra chromosome (number 21) characterized by a typical physical appearance and mental handicap.

Etiology:

1. An error in cell division
2. Associated with advanced maternal age
3. Abnormal attachment of chromosomes inherited from parents

Clinical Manifestations:

1. Eyes slant upward and outward
2. Excess skin on back of neck
3. Flattened nasal bridge and flat facial profile
4. Small ears often incompletely developed
5. Short broad feet
6. small male genitalia
7. Absence of moro reflex

Complications:

1. Recurrent infection of upper respiratory tract
2. Skin infection
3. Serious behavioral problems

Nursing Responsibility:

1. Establishing and maintaining adequate nutrition
2. Recording and reporting any sign of physical complaints
3. Providing a safe environment for the infant
4. Providing appropriate stimulation according to child's age
5. Encouraging parental participation in caring for and handling the infant
6. Working effectively with parents

13.4 Cretinism/ hypothyroidism:

Defects or complete absence of the thyroid gland with insufficient production thyroid hormone causes severe retardation of physical and mental development of the child, sometimes known as cretinism. Early detection and proper treatment can provide a normal life.

Clinical Features:

1. Delayed growth and development
2. Signs of slow metabolism (hair and skin dry, thick and coarse)
3. Goiter may be present or thyroid gland may be absent

Management:

1. Thyroid hormone treatment should be continued for life
2. If diagnosed within the first month of life and substitution with thyroid hormone continued regularly the child will have a normal life
3. With late diagnosis and inadequate treatment the child will be severely retarded and handicapped

13.5 Juvenile Diabetes Mellitus:

The characteristics of juvenile diabetes mellitus differ from those when the onset is later in life.

Symptoms:

The onset can be insidious with loss of appetite, weight loss, abdominal cramps, vomiting, emotional disturbance and lassitude. Polyuria, and polydipsia in young children are symptoms to make you think of diabetes mellitus which is not uncommon and is often overlooked

Findings of sugar in urine, and an increased blood sugar prove the diagnosis.

Treatment:

1. Insulin injection must be given for life
2. Lengthy hospital admission is often necessary
3. If a child is comatose and the test for sugar in the urine is strongly positive
 - set IV drip with normal saline
 - give 1u/kg crystalline insulin im
 - refer the child urgently to hospital

Prognosis:

The prognosis is poor because the management is difficult and complications like hypoglycemia and vascular changes occur frequently.



Bibliography:

1. Addele Pilliteri (1987), Child health Nursing, care of the growing child, Little, Brown and Company.
2. Addele Pilliteri (1992), Maternal and child health Nursing, care of the child bearing and child rearing family, J.B. Lippincott Company.
3. Adele Pilliteri child health nursing, care of the growing family, 3rd edition
4. AMRF (1975), Child Health, a manual for Medical assistants and other rural health workers, Nairobi Kenya.
5. AMRF (1984) Child Health Manual for Medical assistant and other health workers Nairobi, Kenya.
6. AMRF, (1984), Manual for rural health workers, Nairobi Kenya AMRF (1975), Child Health, a manual for Medical assistants and other rural health workers, Nairobi Kenya.
7. Behrman Kliegman. Essentials of paediatrics, 2nd edition.
8. Child Health Manual for Medical assistant and other health workers, by AMRF, 1984.
9. Dickinson Silver man, maternal and infant nursing care. 2nd edition Mosby Champanny
10. Dorothy R.Marlow, R.N. (1977), Text Book of paediatric nursing, W.B Saunders Company, fifth eddition
11. E.B.Turya, F.A Tiu (1991), Essential Paediatric Nursing, Chelmstord, Kampala

12. EPI models
13. EPI National strategy (If you can get access from MOH)
14. GJ. Ebrahim (1993) Pediatrics practice in developing countries, 2nd edition.
15. Katharyn May, Laura Mahlmeister (1990), Comprehensive Maternity Nursing, nursing Process and the child bearing family, J.B.Lippincott Company, second eddition.
16. Lillian S. brunner and Doris S.Suddarth (1988), Manual of Paediatric Nursing, second eddition.
17. Manual on maternal and child care, MOH, Addis Ababa, Ethiopia 1995.
18. Mark H. Swartz (1989), Text book of physical diagnosis, W.B Sauders Company.
19. Marlow, textbook of paediatrics nursing, 5th edition. Philadelphia condom 1997
20. Maurice King, Felicity king (1978), Primary Child Care, a manual for health workers, Oxford University Press.
21. MOH (1995), Manual on Maternal and Child Health Care, Addis Ababa, Ethiopia.
22. Paediatrics practice in developing countries, GJ. Ebrahim 2nd edition 1993.
23. Patricia Wiel and Cadewing, essentials of maternal and Newborn Nursing 2nd edition.
24. WHO (1995) Treatment and prevention of acute diarrhea, guide lines for the trainers of health workers. Geneva,