

# MODULE



Funded under USAID Cooperative Agreement No. 663-A-00-00-0358-00.

Produced in collaboration with the Ethiopia Public Health Training Initiative, The Carter Center, the Ethiopia Ministry of Health, and the Ethiopia Ministry of Education.

### **Important Guidelines for Printing and Photocopying**

Limited permission is granted free of charge to print or photocopy all pages of this publication for educational, not-for-profit use by health care workers, students or faculty. All copies must retain all author credits and copyright notices included in the original document. Under no circumstances is it permissible to sell or distribute on a commercial basis, or to claim authorship of, copies of material reproduced from this publication.

**©2005 by Tefera Belachew, Abebe Gebremariam, Worku Legesse,  
Tsegaye Asres, Seid Tiku, Estifanos Kebede, Tamirat Tigabu,  
Getachew Dememe, Muluneh Haile**

All rights reserved. Except as expressly provided above, no part of this publication may be reproduced or transmitted in any form or by any means, electronic or mechanical, including photocopying, recording, or by any information storage and retrieval system, without written permission of the author or authors.

*This material is intended for educational use only by practicing health care workers or students and faculty in a health care field.*

## AKNOWLEDGEMENTS

The development of this module has gone through series of individual and group works, meetings, discussions, writings and revisions. We would like to express our deep appreciation to The Carter Center, Atlanta Georgia, for their financial support of the activities in the development of this module all the way through.

The input of many academicians at home and abroad has contributed a lot to the development of this module. We highly appreciate the inputs of international reviewers( Dr. Troy A Jacobs and Dr. Elizabeth Downes) in making this module to assume its current shape.

We also like to extend our thanks to the Faculty of Public Health, Faculty of Medical Sciences, Population and Family Health Department, Health

# TABLE OF CONTENTS

<b>ACKNOWLEDGEMENT</b> .....	i
<b>TABLE OF CONTENTS</b> .....	ii
<b>UNIT ONE: Introduction</b> .....	1
1.1 Purpose and use of the module .....	1
1.2 Directions for using the module .....	1
<b>UNIT TWO: Core modules</b> .....	3
2.1 pre and posttests .....	3
2.2 Significance and brief description of the problem .....	19
2.3 Learning objectives .....	25
2.4 Case study: learning activities .....	27
2.5 Definition .....	30
2.6 Epidemiology .....	31
2.7 Etiology and pathogenesis .....	37
2.8 Clinical features (signs and symptoms) .....	39
2.9 Prevention of selected micronutrient deficiencies .....	40
2.10 Learning activities (case study) continued .....	43
<b>UNIT THREE: Satellite Modules</b> .....	44
3.1. Health Officer .....	45
3.2 Public Health Nurse .....	63
3.3 Medical Laboratory Technician .....	88
3.4 Sanitarian .....	102
3.5 Primary health worker/ Community health worker .....	113
3.6 Take home message for care givers .....	121
<b>UNIT FOUR: Role and task analysis</b> .....	124
<b>UNIT FIVE: Glossary and abbreviations</b> .....	139
<b>UNIT SIX: References</b> .....	142
<b>UNIT SEVEN: Annexes</b> .....	153
7.1. Answer keys .....	153
7.2. Other materials .....	158

# UNIT ONE

## INTRODUCTION

### 1.1. PURPOSE OF THE MODULE

The lack of appropriate and relevant teaching materials is one of the bottlenecks that hinders training of effective, competent task oriented professionals who are well versed with the knowledge, skills, attitudes and that would enable them to solve community problems. Preparation of such a teaching material is an important milestone in an effort towards achieving these long-term goals.. Therefore, this module is prepared for equipping trainees with adequate knowledge, skills and attitudes through interactive teaching mainly focused on Micronutrient deficiency.

The preparation of this module has taken in to account the current guideline on the management of severe acute malnutrition, guideline on infant and young child feeding, the essential nutrition actions approach and guideline on micronutrient deficiency prevention and control of the Federal democratic Republic of Ethiopian Ministry of health.


This module can be used for the basic training of health center teams at the training institutions and training of health center teams who are already in service sectors, and by health extension workers and care givers. However, it was not meant to replace standard textbooks or reference materials but can be used as a supplement to enhance the teaching and learning process.

### 1.2 DIRECTION FOR USING THE MODULE

In order to make maximum use of the module the user should follow these directions:

- 1.1.1 Check prerequisite knowledge required to use the module
- 1.1.2 Do the pretest pertaining to the core module in section 2.1.1
- 1.1.3 Read the core module thoroughly.

- 1.1.4 After going through the core module try to answer the pretest questions.
- 1.1.5 Evaluate yourself by referring to the key given in section 7.1 and 7.2
- 1.1.6 Read the case study and try to answer questions
- 1.1.7 Use the listed references and suggested reading materials to substantiate and supplement your understanding of the problem
- 1.1.8 Look at the satellite module and the task analysis related to your field to understand your role in the team in managing a micronutrient deficiency.



## UNIT TWO

### CORE MODULES

#### 2.1. PRE-AND POST TEST

##### 2.1.1 PRE-AND POST TEST FOR THE HEALTH CENTER TEAM (FROM THE CORE MODULE)

**Directions: Choose the letter of the choice with the right answer.**

1. Which segment of the population is more predisposed to Iodine deficiency disorders?
  - a. Under two years
  - b. Pregnant women
  - c. Adults
  - d. Adolescents
  - e. A and B
  
2. What are the different risk factors involved for the development of IDD?
  - a. Ignorance about the importance of iodized salt
  - b. Poor consumption of seafood like fish
  - c. Increased consumption of food that contains thiocyanate or cyanogenic glycoside
  - d. All of the above
  - e. A and C
  
3. Which of the following is not true about IDD?
  - a. Total goiter rate is 26% in Ethiopia
  - b. IDD is more common in mountainous areas
  - c. IDD affects also other mammals
  - d. Can be prevented by fortification of foods and drinks
  - e. None
  
4. List 4 consequences of IDD?

- a. ....
- b. ....
- c. ....
- d. ....

5. What are the basic causes of anemia?

- a. Inadequate dietary intake
- b. Excessive RBC destruction
- c. Inadequate RBC production
- d. All of the above
- e. None

6. Which of the following is a false statement?

- a. IDD is associated with the use of food like cassava
- b. Immunization can prevent IDD
- c. Xerophthalmia is more prevalent among children below 6 years
- d. IDD is non-preventable communicable disease
- e. b and d

7. What preventive measures should be taken to prevent VIT A deficiency?

- a. Diarrhea control using ORS
- b. Fortification of food with vitamin A
- c. Immunization program to control measles
- d. All of the above
- e. None of the above

8. In the clinical work up of anemia what laboratory investigations can be done in a routine laboratory setup?

- a. Hemoglobin determination
- b. Stained red blood cell morphology assessment
- c. Stool examination for ova of parasites




- d. Reticulocyte count
- e. All of the above

9. What is the importance of reticulocyte count in the assessment of Anemia?

- a. to diagnose anemia
- b. to diagnose polycythemia
- c. to assess the presence of abnormal red blood cell morphology
- d. to assess the status to RBC production with in the bone marrow

9. What is the importance of reticulocyte count in the assessment of Anemia?



- a. Calcitonin
- b. Parathyroid hormone
- c. 1,25 dihydroxycholecalciferol
- d. All
- e. None

14. Which of the following is not the function of zinc?

- a. Promotes growth
- b. Enhances resistance to infection
- c. Acts as antioxidant
- d. Forms part of insulin
- e. None

15. Which of the following is not a consequence of a zinc deficient diet during pregnancy?

- a. Delivery of low birth weight baby
- b. Pre-term labor
- c. Precipitated labor
- d. Post partum hemorrhage
- e. None

16. Which of the following foods are **excellent** sources of zinc?

- a. Dairy products
- b. Dark red beef
- c. Green leafy vegetables
- d. Cereals
- e. Legumes

17. One of the following is not the consequence of folate deficiency?

- a. Megaloblastic anemia
- b. Spinal cord and peripheral neural disease
- c. Aggravates immunity in HIV infected patients
- d. Neural tube defect during fetal period

e. None

18. Vitamin A is useful for the following body functions?

- a. Epithelial tissue replication
- b. Bone growth
- c. Vision
- d. Reproduction
- e. All

19. Which of the following vitamins cause hypervitaminosis?

- a. Vitamin B-9 (folate)
- b. Vitamin A
- c. Ascorbic acid (vitamin C)
- d. Vitamin D
- e. b and d

20. One of the following is not true about folic acid?

- a. Found in good amount in liver, kidney, and fish
- b. Important for the synthesis of hemoglobin during RBC synthesis
- c. Needed most during growth and pregnancy
- d. Important for myelin sheath synthesis
- e. Its' deficiency causes pernicious anemia

21. Which one of the following is the clinical manifestations of rickets?

- a. Frontal bossing
- b. Double malleoli
- c. Delayed fontanelle closure
- d. Bow legs
- e. All

22. One the following deficiencies causes megaloblastic anemia: -

- a. Vit D deficiency

- b. Iron deficiency anemia
- c. Folic acid deficiency
- d. Vit B-12 deficiency
- e. c and d

23. Which of the following is the correct order of vitamin A deficiency clinical manifestations?

- a. Night blindness, Bitot's spot, corneal xerosis, corneal ulceration, corneal scar
- b. Bitot's spot, corneal xerosis, night blindness, corneal scar, corneal ulceration
- c. Night blindness, corneal xerosis, corneal ulceration, Bitot's spot, corneal scar
- d. Bitot's spot, night blindness, corneal xerosis, corneal ulceration, corneal scar
- e. Corneal xerosis, night blindness, Bitot's spot, corneal ulceration, corneal scar

24. Osteoporosis is not caused by:

- a. Vit B 1 deficiency
- b. Vit D deficiency
- c. Steroid treatment
- d. Calcium deficiency
- e. Menopause

25. Which of the following nutrients are not anti-oxidants?

- a. Vitamin A
- b. Vitamin E
- c. Vitamin C
- d. Zinc
- e. Iron

26. Which of the following nutrients are suitable for fortification?

- a. Vitamin A

- b. Vitamin D
- c. Vitamin B2 (Riboflavin)
- d. Iodine
- e. All

27. Which of the following nutrients affects bone integrity?

- a. Vitamin A
- b. Vitamin C
- c. Vitamin D
- d. Calcium
- e. All

28. One of the following causes of anemia does not match the type anemia listed?

- a. Microcytic hypochromic anemia – iron deficiency
- b. Microcytic hypochromic anemia – Hookworm infection
- c. Macrocytic anemia – Folate deficiency
- d. Normocytic normochromic anemia – Bone marrow failure
- e. Pernicious anemia – *Diphyllobotrium latum* infection

### 2.1.2. PRE AND POSTTEST FOR SPECIFIC CATEGORIES OF THE HEALTH CENTER TEAM (FROM THE SATELLITE MODULES)

#### 2.1.2.1. PRE AND POST TEST FOR HEALTH OFFICERS

**Directions:** - Circle on any of the following choices that you think is correct answers.

1. One of the following is a nutritional problem of public health importance in developing countries,
  - a. Protein energy malnutrition
  - b. Iron deficiency anemia (IDA)
  - c. Iodine deficiency disease (IDD)
  - d. Vitamin A deficiency (VAD)

e. Vitamin D deficiency

2. Which groups of the population are more predisposed to IDD?

- a. Under two years of age
- b. Pregnant ladies
- c. Adults
- d. Adolescents
- e. A and B

3. What are the different risk factors involved in the development Iodine deficiency disorder?

- a. Low socioeconomic conditions
- b. Ignorance of about the importance of iodized salt
- c. Infections like measles, Pertussis, diarrhea
- d. Use of some drugs
- e. All of the above

4. How prevalent is IDD in Ethiopia?

5. List 3 different types of health problems encountered because of IDD?

- a. ....
- b. ....
- c. ....

6. What are the basic causes of anemia?

- a. Inadequate dietary intake
- b. Excessive RBC destruction
- c. Inadequate RBC production
- d. All of the above
- e. None

7. Which of the following is a false statement?

- a. IDD is associated with the use of food like cassava
- b. Immunization can prevent IDD
- c. Exophthalmia is more prevalent among children below 6 years
- d. IDD is a non-preventable communicable disease
- e. B and D

8. What preventive measures should be taken to prevent VIT A Deficiency

- a. Diarrhea control using ORS
- b. Fortification of food with vitamin A.
- c. Immunization program to control measles
- d. All of the above
- e. None of the above

9. In the clinical work up of anemia, what laboratory investigations can be done in a routine laboratory setup?

- b. It enables the classification of anemia
- c. To diagnose iron deficiency anemia
- d. All of the above

12. What pathogens contribute indirectly to the development of anemia?

- a. Viruses
- b. Bacteria
- c. Parasites
- d. All of the above

13. The basic objective of managing a child with Vit A deficiency: -

- a. Treating superimposed infections
- b. Supplying therapeutic dose of Vit A
- c. Managing complications
- d. Provision of immunization (Vit A)

14. The commonest type of malnutrition in Ethiopia

- a. Iodine deficiency disorders
- b. Vitamin A deficiency
- c. Iron deficiency anemia
- d. Rickets
- e. None

15. Indication for atherapeutic dose of vitamin A is a case of

- a. Xerophthalmia
- b. Measles
- c. Severe PEM
- d. Severe ARI



e. A and B

16. Which of the following diseases have a very close relationship with protein energy malnutrition

- a. Tuberculosis
- b. Measles
- c. Diarrhea
- d. Pertusis (whooping cough)
- e. Common cold

Abebech brought her 3 years old male child called Temtime to the pediatric OPD of Jimma Hospital. She told you that the child has intermittent diarrhea, loss of appetite. Besides she stated that the child is not interested in his surrounding and sits miserably. On physical examination you found out that the child is apathetic, hypotensive, has gray easily pluckable hair, edema, weighs 9kg when he is expected to weigh 14kg. He also has bowed legs and bead like swellings on the costochondral junctions. The child is also pale and has difficulty of vision during dim light and had whitish triangular patches on the lateral lower quadrant of the bulbar conjunctiva. Answer questions 5 to 10 based on the above scenario.

17. How would you manage this child?

- a. Give him cod liver oil and advise the mother to expose him to sunlight
- b. Start him with low protein 1-1.5g/kg/d and low energy 100 kcal/kg/d in the stabilization phase and later increase to 5 gm of protein/kg/d and 180kcal/kg/d in the rehabilitation phase
- c. Give iron after 7 days (after control of infection)
- d. 200,000 IU of vitamin A is required at the 1<sup>st</sup>, 2<sup>nd</sup> and 14<sup>th</sup> days
- e. All are correct

18. What will be your approach to the mother to prevent a recurrence of the situation?

- a. Nutrition education on child feeding and meal planning
- b. Counseling her on the importance of mixing different foods cereals, legumes, animal products, sea foods; fruits and vegetables in child feeding to prevent micronutrient deficiency
- c. Tell her the importance of gardening in her yard-garden if she has land
- d. Work with her how to improve the nutritional status of her child and appoint her for follow up (growth monitoring)
- e. All are correct

19. Other micronutrient deficiencies that co-exist with PEM include:

- a) Vitamin A deficiency
- b) Vitamin D deficiency
- c) Riboflavin deficiency
- d) Iron deficiency

#### 1.2.1.2. PRE AND POST TEST TO PUBLIC HEALTH NURSES

**Direction: Respond to the following questions accordingly.**

b36(Lco-edowher once nuciticontimpor)]TJ-0.0004 T4 0.11763.2612.325 0acornd appiher onhs cst c(

- D. \_\_\_\_\_
- E. \_\_\_\_\_
- F. \_\_\_\_\_

2. Who are at the risk of developing Vitamin A Deficiency?

- A. Children at the age of six months and six years.
- B. Pregnant women
- C. Lactating mother
- D. All of the above.

3. List down the three methods used to control vitamin A deficiency?

- A. \_\_\_\_\_
- B. \_\_\_\_\_
- C. \_\_\_\_\_

4. Who are at the risk of developing Iodine deficiency diseases?

- A. School age children
- B. women of reproductive age
- C. Pregnant and lactating mothers
- D. All of the above.

5. Who are at the risk of developing Iron deficiency Anemia?

- A. Non Lactating mother.
- B. Exclusively breast fed infant.
- C. Pregnant women and children.
- D. The answer is not given.

6. Which one of the following is the public health measure to control iron deficiency anemia?

- A. Iron Supplementation
- B. Fortifications of food.
- C. De-worming

D. All of the above.

7. What is the recommended daily requirement for iodine per person to prevent IDD (Iodine Deficiency Disease)?

8. Which one of the following vitamin enhances the absorption of Iron?

- A..Vitamin A
- B. Vitamin B
- C. Vitamin D
- D. Vitamin C

9. List some of food items that are of animal origin and that improves iron availability in the body?

---

---

---

---

10. List at least five causes of iron deficiency anemia?

- A. ....
- B. ....
- C. ....
- D. ....
- E. ....

11. What does the term "Xerophtalmia` mean"?

**2.1.2.3. PRE AND POST TEST FOR MEDICAL LABORATORY TECHNICIANS**

#### 1.2.1.4. PRE AND POST TEST ON PEM FOR THE SANITARIANS

**Direction.** Circle on any of the following choices which you think is the best answer.

1. Asebech always wraps her eight months old child with cloths and the child had no a chance to be exposed to sunlight for fear of cold and evil eyes. As a result the child's legs were weak and not straight. What will be the possible cause(s) of the problem?
  - A. Lack of fish intake
  - B. Evil eyes
  - C. Cold
  - D. Inadequate sun light resulting in vitamin D deficiency
2. What type of environmental health intervention shall be implemented in areas where iron deficiency anemia is highly prevalent?
  - A. Proper human waste management
  - B. Food hygiene
  - C. Provision of safe water supply
  - D. All
3. Animut, a fisherman who is living in Gorgora at the edge of Lake Tana, was visiting Gorgora clinic with a complaint of body weakness. The physical examination and laboratory test revealed that Animut's problem was anemia. What will be the possible cause(s) of anemia based on the explanation given above?
  - A. Lack of fish intake
  - B. High fish intake
  - C. Schistosomiasis
  - D. Lack of *Teff*
4. Inhabitants of Kitimbile Pesant association in Jimma zone lead a subsistence life where it is to buy vegetables and animal products. Which deficiency disease is likely to occur?
  - A. Anemia
  - B. Night blindness
  - C. Rickets
  - D. Marasmus

### 2.1.2.5. PRETEST FOR HEALTH EXTENSION WORKERS (HEWS)

1. The causes of micronutrient deficiency is (are)
  - a. Germs
  - b. Evil eye
  - c. Not proper child feeding practices
  - d. Tooth extractions
  - e. None of the above
  
2. One of the following is not a method of preventing iron deficiency anemia: -
  - a. Maintaining personal hygiene and proper waste disposal
  - b. G wastTT1 1 T02 Tc T\*[(c. )160o1x waste dislm prmr dN m .735ically

- b. Corneal scar
  - c. Growth retardation
  - d. Poor school performance
  - e. Susceptibility to Infection
6. Folic acid deficiency causes all of the following except,
- a. Night blindness
  - b. Neural tube defect during pregnancy
  - c. Neuropathy in adults
  - d. Anemia (Megaloblastic)
  - e. None

## **2.2. SIGNIFICANCE AND BRIEF DESCRIPTION OF THE PROBLEM**

### **2.2.1. IODINE DEFICIENCY DISORDER**

Iodine deficiency disorder refers to medical conditions ranging from simple goiter to deafness, mutism, squint and profound mental retardation called cretinism. Iodine is part of the hormone thyroxin, which controls energy metabolism. The deficiency diseases are goiter and cretinism. Iodine occurs in seafood and in foods grown on land that was once covered by oceans. The daily requirement for a normal person is 150 micrograms. The term iodine deficiency disorders' (IDD) has been adopted to describe the spectrum of effects of iodine deficiency that include goiter, endemic cretinism psychomotor delays, and subsequent increased pre-and post-natal mortality.

Universal salt iodization (USI) is the most widely practiced intervention in eliminating iodine deficiency disorders (IDDs). Salt iodine test is an imprecise method to describe the prevalence of IDD. Salt iodine test is an imprecise method to describe the prevalence of IDD.





bone. However, its clinical significance is mainly due to the effect the disease has on general child health. Developmental delay is a commonly observed finding in rickets. Moreover, rickets is associated with increased morbidity and mortality. There is ample evidence demonstrating increased rates of diarrhoeal diseases and acute respiratory infections, in rickets. An increased susceptibility to infection associated with rickets may be multi-factorial: the chest wall deformity of late rickets alters immune status and the association of rickets with Protein-energy malnutrition (PEM) are some of the factors involved.

Vitamin D deficiency is the commonest cause of rickets in children and vitamin D supplementation has brought down the incidence of rickets to a virtually non-existent level in the developed world. The infant with rickets in the United States, for instance, is typically a black breast-fed infant who does not receive supplemental vitamin D and usually lives in northern areas of United States with lower levels of sunlight. In developing countries where there is a high proportion of sunny weather, the occurrence of rickets in a significant proportion of children is largely attributed to social and cultural factors. Keeping children indoors because of the fear that they will catch cold, fear of evil eyes, and the tradition of wrapping children in swaddling clothes contribute to lack of adequate sunlight exposure. Calcium administration is appropriate in all cases of rickets. The preventive measures for rickets may also need to be modified. Calcium supplementation along with sunlight exposure is a more effective strategy.

#### **2.2.4. VITAMIN A DEFICIENCY AND XEROPHTHALMIA**

Vitamin A deficiency is one of the major nutritional problems in Ethiopia as well as many developing countries. In spite of the potential to produce a wide variety of fruits and vegetables containing beta-carotene, most people do not consume them due to the reason that they tend to sell them (E.g. Carrots, mangos, Papayas) on the one hand and the traditional belief that these foods are low class foods (Kale, Cabbage). As a result mostly children under the age of 59 months are affected by the tolls of morbidity, disability and mortality due vitamin A deficiency. Severe vitamin A deficiency (VAD) has been known to cause xerophthalmia (X). Xerophthalmia (X) literally means “dry eye” in Greek.

Vitamin A deficiency is considered to be a problem of public health importance if one of the following is observed in children under five of that area (Table1)

*Table 1. Threshold for making VAD a problem of public health Importance*

INDICATOR	LEVEL	PREVALENCE OF VAD IN UNDER FIVE CHILDREN THRESHOLD (%)
Xerophthalmia	Night blindness (XN)	1.0
	Bitot's spot (XIB)	0.5
	Corneal xerosis(X2) or corneal ulceration/ Keratomalacia < 1/3 of corneal surface(x3A), or corneal	

Short study of children with/without non-corneal xerophthalmia in Indonesia (Sommer et al 1983)

Type of Xerophthalmia	Relative Risk of dying in next 3 months
XN night blindness only	2.4
XIB (Bitot's spot only)*	6.6
Both Night blindness and bitot's spots (XN and XIB)	8.6

\* See glossary

Risk in children with xerophthalmia divided by risk in children without xerophthalmia. Clinical vitamin A deficiency is strongly associated with subsequent mortality in young children. Vitamin A supplementation increases child survival by decreasing the severity of diseases like measles, acute respiratory infections, diarrheal disease and acute malnutrition. It decreases mortality from measles by 40%, mortality from all the above common infections by 23-35%. Vertical transmission of HIV from mother to child is also lower in mothers with normal vitamin A status.

### 2.2.5. ZINC

Zinc is found in every cell of the body and is a part of over 200 enzymes. Essential for the maintenance of vision, taste and smell, this mineral is also necessary for immune function, protein synthesis, and cell growth. It is required for the activity of the antioxidant enzyme super oxide dismutase. Zinc has been used successfully in the treatment of rheumatoid arthritis, acne, and macular eye degeneration.

While severe deficiencies of zinc are uncommon, marginal deficiencies are frequent in the elderly, those suffering from abnormal eating behavior, and AIDS patients. Deficiencies may be responsible for many of the secondary conditions in AIDS patients, including gastrointestinal malfunction, diarrhea, anorexia, impaired immunity, central nervous system malfunction, and hypoalbuminemia.

Zinc plays an important role in immune function, helping increase resistance to infection. When combined with Vitamin A supplements, zinc helps the suppressed immune system recover. Low levels of zinc result in a decrease in helper T-cells and thymic hormone, which may adversely effect immune functioning. Low doses of zinc have been shown to improve immunity without causing the side effects associated with high doses of the element. It has shown antiviral properties and is used in lozenge form to shorten the duration of the common cold. In stark contrast, high doses of zinc may also impair immunity. Zinc is found from red meet, the darker the meat the more its zinc content is. Optimal level of zinc is 15-30 mg.

#### **2.2.6. FOLIC ACID**

##### **B9)**

Vitamin B-9, more commonly known as folic acid, functions together with a group of related water-soluble compounds, collectively called folacin, in many body processes. It is critical to cellular division because it is necessary in DNA synthesis.

Folic acid, also known as folacin or folate

Due to the overwhelming evidence of folacin's impact on preventing neural tube defects, the FDA is now recommending food fortification of this vitamin. It found in green leafy vegetables cereals and legumes.

### **2.3. LEARNING OBJECTIVES: -**

For effective prevention and management of a case with micronutrient deficiencies the students at the end of the training will have the following knowledge, attitudes and behavioral outcomes: -

1. Define and identify the types of micronutrient deficiencies
2. Enumerate the causes and factors contributing to micronutrient deficiencies
3. Describe the magnitude and contribution of micronutrient deficiencies to the overall child health problems in the country and locally.
4. Identify and describe the clinical manifestations of various micronutrient deficiencies and its complications.
5. Demonstrate the process of assessing a child with micronutrient deficiencies
6. Identify the degree of micronutrient deficiencies
7. List the diagnostic methods and procedures for a case with micronutrient deficiencies.
8. Describe the principles and methods of treating micronutrient deficiencies
9. Select the appropriate treatment for a case of micronutrient deficiencies
10. Describe methods of preparing dietary treatment for a case of micronutrient deficiencies.
11. Identify and manage or refer timely when needed, a case of severe micronutrient deficiencies
12. Demonstrate the appropriate management of a case with micronutrient deficiencies
13. Learn how to monitor their growth (growth monitoring) and take action

14. Promote hygienic preparation and storage of weaning foods.
15. Identify methods and targets for health education in the prevention of micronutrient deficiencies
16. Describe proper growth monitoring activities and their importance in the prevention of micronutrient deficiencies
17. Promote breast feeding and proper weaning practice
18. Promote immunization of children



## 2.4. CASE STUDY: A LEARNING ACTIVITY

**Case study 1.** Micronutrients can be plenty out there but inhibitory prices make them Unavailable to the consumers!



In a large city was a poor neighborhood called Village X. There was no access to fresh produce so people bought all their food in markets and stores but food prices were so high that families were never able to buy enough. The main cause of the high prices was the middlemen. The middlemen were people between the farmers in the country who grew the food and the families in Village X who ate it.

Maize and *teff* meal go to the families of Village X. Out in the country, farmers grow maize and *teff* in their fields. After the maize and *teff* was harvested they put them in sacks. A young man bought the sacks of maize and *teff* and took them to a small town. A man at the store bought the sacks and kept them in his grain store. A truck driver picked up the sacks and took them to a mill. The mill owner took the maize and *teff* into flour and put it back into sacks. Another truck driver took the sacks to big markets. A young man bought a sack of maize and *teff* flour and took it to Village X. There he opened the sack and sold the meal to six market women. The market women took the maize and *teff* to the Village X market for selling.

The middlemen between the farmwomen and the Village X mothers were middlemen. As the maize and *teff* passed from one person to another the price went up. The price of *teff* was roughly four times that of maize. The Village X people were

too poor to pay for *teff* meal or meat products, they resorted to buying maize every time because it was relatively cheaper. An Epidemiological survey in Village X revealed many residents with symptoms of loss of appetite, pallor, lassitude, weakness and children in that area were poorly attentive at schools and scored lower grades. **The problem was that mothers could afford to buy only maize which is deficient in iron leading to the development of a deficiency disease known as anemia.** This problem could be easily corrected by reducing the price of *teff* to an affordable level, which may involve controlling the middlemen by concerned authorities.

### Case study 2: Night blindness



Everyday the night falls Adanech experienced an inability to see clearly. Her uneducated father and mother were anxious about her situation and were trying to improve her sight and took her to almost all religious settings, including an exorcism in the area. However, her sight was getting worse day by day and reached a point where she needed help to walk every time after a sunset. Village girls now labeled her as “*daphintam*”- meaning unable to see at night. As the youngest girl in a family of 10, what went wrong to her sight?

Adanech was born in a rural village located 30 km North of Addis Ababa. All her brothers and sisters before her grew up in a time where there was huge expanse of farm and grazing land and hence fruits, and dairy products were plentiful. There was always a surplus of fruits and animal products for sale to city dwellers and in return to buy salt, oil, and gasoline and other household items that are no







and more than 150 million people are affected in Africa alone. In Ethiopia, one out of every 1000 is **a cretin** (severely mentally retarded dwarf), and about 50,000 peri-natal deaths are occurring annually due to IDD. Twenty two percent of the total population have goiter and 62 % of the population is at risk for IDD according to national survey conducted by the previous Ethiopian Nutrition Institute. In some pocket areas of the country the prevalence of goiter is found to be 50-95 % (WHO considers that if the goiter rate is above 5 % in the population it is a public health problem). From the various surveys conducted in many parts of Ethiopia, IDD has become one of the biggest nutritional public health concerns. Various surveys of goiter prevalence showed a particular predilection to the highlands and inland areas with the prevalence ranging from 0.4% to 63%.

## 2.6.2. IRON DEFICIENCY ANEMIA

Nutritional problems are common

Ascorbic acid (Vitamin C) is one of the factors in the diet that enhances absorption of non-heme iron. The influence of ascorbic acid is most substantial in inhibitory food, which means food that contains the main inhibitors of nonheme iron absorption, phytate and polyphenoles. It is effective in promoting iron absorption if eaten together with the iron. There could be a decrease in the consumption of sources of this vitamin either due to seasonally availability, economical problems or lack knowledge.

Organic acids also have similar role in promoting iron absorption. It was shown that the addition of citric, malic or tartaric acids to rice-based meal highly improved iron absorption. These organic acids are obtained from vegetables. Fermentation of foods also improves the bioavailability of non-heme iron. Demant et al 1980-noted increased absorption of iron from South-African beer made from





Rickets also contributes to increased morbidity, disability and mortality. In a case-control study done in Addis Ababa the odds of dying in rachitic patients was five times higher than that of controls.

Paradoxically, rickets is widely prevalent in many tropical and subtropical regions despite abundant sunshine. Reports from health institutions suggest that rickets is an important child health problem in Ethiopia. In a review of pediatric admissions in Jimma hospital about 7% of under 5 children were diagnosed with rickets. A more recent community based study in Jimma town showed rickets prevalence of 4% in randomly selected children between 6-59 months of age.

The incidence of rickets is particularly high in slum children who live in crowded houses almost devoid of sunlight. Vitamin D deficiency is the most common cause of rickets in Ethiopia. All the rachitic children were said to be exposed to sunlight, the majority daily, and with no clothes on. From clinical experience too some children present with rickets despite history of adequate sunlight exposure (unpublished observation. The deficiency of calcium may increase the requirement for vitamin D, and therefore calcium deficiency may aggravate a borderline vitamin D status. In rats the rate of inactivation of 25-hydroxy-cholecalciferol in the liver is increased by calcium deprivation (or increased phytic acid). The effect is mediated by 1,25 dihydroxy-cholecalciferol produced in response to secondary hyperpara-thyroidism. Studies conducted in Nigeria suggest that rickets is

recommended daily allowance of 400 micrograms (for pregnant women) and 180 mcg for non pregnant women 15 years and above.

Evidence indicates that the problem of folate deficiency is common in the developed







**Emotional deprivation-** In orphan children and in children whose parents are negligent in giving care to their children for different reasons, children will lose their appetite and hence end up in state of malnutrition

**The Cultural factors-** Different biases as to who should take the lion's share of the family 's food (Example, age bias—older children are given more food than the smaller ones,

**Gender bias**—male children are more favored in getting nutritious food than female children in some families, etc.)

**Mal-distribution of food stuffs** - within the family, it occurs between the different ages and sexes due to biases, food prejudices and taboos. It also occurs between the different regions of any country because of inappropriate food and nutrition policy, poor marketing and distribution system due to different reasons like embargo, country under-siege, etc.

## 2.7.2. ETIOLOGY OF MICRONUTRIENT DEFICENCIES

### 2.7.2.1. IDD

Living in the mountainous areas where plant foods have inadequate in iodine

Taking cassava tuber as a food without detoxifying it and other chelating substances

#### 2.7.2.4. Iron deficiency anemia



Cultural malpractice and beliefs in child feeding and weaning process, weaning foods, exposure of children to sun light, time of weaning and food prejudices

Intra household mal-distribution of food (age and sex bias)

Effects of emotional deprivation and neglect on nutritional status of children and proper child treatment practices

Importance of breast feeding

Hygiene (personal hygiene, food hygiene, environmental hygiene)

Importance of immunization

Importance of growing fruits and vegetables in the backyard garden and consumption by the household members regardless of their age and sex.

Parents / caregivers need to be instructed how to modify the nutrient contents of the locally available foods used in weaning and child feeding.

### 2.9.2. Dietary modification: -

This approach focuses on modifying the micronutrient y, protein and energy content of the complementary foods.

- Making iodized accsssible to the whole family especially women in child bearing age.
- Iron and folic acid supplementation to mother during pregnancy and lactation
- vitamin A supplementation to the mother during lactation during the first 6 weeks of delivery
- Iron and vitamin A supplementation to children
- Periodic deworming(giving antihelmithic ) of mothers( except in the first trimester of pregnancy) and children every 6 months

### 2.9.3 Economic approach:

This approach aims at improving the incomes of the target community as a solution to their nutritional problems. It is considered usually in areas where there are many poor people and if their purchasing power is low as in the case of urban slums and people displaced because of war and other natural calamities. There are different methods in this approach: -

▶ **Food for work**— This involves offering of some work for the poor people and paying them off in terms of food. It is good in that it offsets seasonality in the dietary intake, but it is donor dependent

**Food subsidy** --- This involves subsidizing of either producers or consumers of food by the government. Structural adjustment policies interfere with the materialization of this approach

**Income generating projects**---This method operates in some regions of Ethiopia and involves development of income generating projects in the community to make them

generate fund for buying food. It includes organizing the community and using their potentials in the running of the project. The projects could be weaving, pottery, Bee keeping, etc. This approach needs a good feasibility study on how the income generated is used, the sustainability of the programme, etc.

The above approaches could be used either simultaneously where it applies or independently. This should be determined by doing a thorough strength, weakness, opportunities and constraints (SWOC) analysis.

#### **2.9.4. Nutritional Surveillance**

Targets for surveillance:- Infant & child growth monitoring and promotion(GMP) activities need to be carried out in an integrated manner with other PHC services. Missed opportunities for GM should be fully utilized in such a way that children coming to the health institutions for other purposes are covered in the growth monitoring and promotion (GMP) activities. Besides every child should be regularly monitored for growth performance (growth take up) every month. **Triple A cycle** (assessment, analysis and action) be employed in effecting GM activities.

*Assessment includes regular measurement of weight and heights of < 5 children and comparing their growth performance with the standard(determining the direction of growth of the child)*

*Analysis includes assessment of the different causes of growth faltering and options for intervention with the mother/care giver.*

*Action involves nutritional intervention to curb the problems, which include counseling of the care taker on child feeding practices.*

The action may include rehabilitation of severely malnourished children and following them up and micronutrient supplementation, Nutrition education on importance of backyard gardening & horticultural activities, dietary diversification, breast feeding and proper child feeding practices.

#### **2.9.5. Nutritional surveys**

Community based nutritional surveys including anthropocentric measurements and dietary consumption surveys need to be carried out among under five children in order to early detect the occurrence of nutritional problems in the community.

## **2.10. LEARNING ACTIVITIES (CASE STUDY) CONTINUED**

Based on the case studies presented in section 2.4, different points of discussion have been incorporated in the respective satellite modules. Therefore, the students are advised to refer to the questions in satellite modules for each professional category and discuss them in the class under the coordination of their facilitator.

## **UNIT THREE**

# SATELLITE MODULES





## 1.2. Direction for using the satellite module

This satellite module can be used in the basic training of Health Center team particularly health officers who are either already in the service or in training Programs. In order to make maximum use of the satellite module, the health officer should follow these directions

Evaluate yourself by doing the pre-test pertinent to your category under section 2.1.2.1 before going through the satellite module and evaluate your self by referring to the answer keys given in the unit 7 section 7.1.1

Read the core module very thoroughly

Read the case study and try to answer questions pertinent to it

Use listed references and suggested reading materials to supplement your understanding of the problem.

For total and comprehensive understanding of the causes, etiology, pathogenesis, epidemiology and prevention of micronutrient deficiencies, the health officer students are advised to refer to the core module.

After going through this module evaluate yourself by doing post-test and comparing your score with the key given in unit 7 section 7.2.1



## **Case study: Learning Activity**

Read the case studies 1,2 and 3 in the core module very thoroughly so that you will be able to answer questions pertaining to it in section 2.12 of this module.

### **2.5. Definition**

*Refer to the core module unit 2 sections 2.5*

### **2.6. Epidemiology**

*Refer to the core module unit 2 sections 2.6*

### **2.7. Cause, Etiology and pathogenesis**

*Refer to unit 2 section 2.7 of the core module*

#### **2.7.2. ETIOLOGY OF MICRONUTRIENT DEFICENCIES**

*Refer to the core module section 2.7*

poor in zinc content

### **2.8. Clinical Features**

#### **2.8.1. Iodine deficiency**

- **Fetus**
  - Abortion
  - Still birth
  - Congenital anomalies
  - Increased infant mortality
  - Neurologic cretinism (including mental deficiency, deaf mutism, spastic diplegia, squint)
  - myxedematous cretinism (dwarfism, mental deficiency)
  - Psychomotor defects

- **Neonate**
  - Neonatal goiter
  - Neonatal hypothyroidism
- **Child and adolocsent**
  - Goiter
  - Juvenile hypothyroidism
  - Impaired mental function
  - Retarded physical development
- **Adult**
  - Goiter with its complications
  - Hypothyroidism
  - Impaired mental function
  - Poor physical work out put
  - Iodine induced hyperthyroidism
- **Beasts of burden (animals used for production like oxen, horses, donkeys are also affected)**
  - Weakness

## Diagnosis

Three components to diagnosing IDD:

1. Determination of thyroid size/goiter
2. Determination of urine iodine excretion
3. Determination of blood T4 +TSH levels

Usually the diagnosis must depend upon clinical assessment only using WHO criteria

## Measuring Thyroid Size /WHO/

STAGE O No goiter

STAGE IA Goiter detectable by palpation and not visible when neck is in the normal position or distended

STAGE IB Goiter palpable and visible when neck is extended

STAGE IIGoiter visible with the neck in normal position

STAGE III Goiter visible at a distance

STAGE IV Huge goiter

At the community level determination of the endemicity of IDD is very important to institute appropriate intervention measures. The following are classification and techniques of survey to detect IDD at the community level.

**Table 1. Classification of Endemia of Iodine deficiency disorders by Severity**

Grades of severity of iodine deficiency disorders			
	Mild IDD	Moderate IDD	Severe IDD
1. Prevalence of goiter (total)	5-19%	20-29%	$\geq 30\%$
2. Cretinism	0	0	0-5
3. Daily urine urinary iodine ug/d	50-100	25-49	$< 25$
Median Urine iodine ug/d	3.5-5	2-3.4	0-1.9
4. Prevalence of neonatal TSH $> 50$ ug/d	$< 1\%$	1-5%	$> 5\%$

Source: WHO/ICCIDD Report

## Survey Techniques for endemic Goiter and Cretinism

1. Survey Organization
  - i. Examine school children
  - ii. If goiter prevalence is  $> 10\%$  in school children, sample population for survey
2. If goiter prevalence is  $> 10\%$  in school children, assess 24 hours urine iodine
3. Check for Goitrogenic factors (e.g. soy products, drugs, cabbage, cassava, etc)

## TREATMENT AND PREVENTION OF IODINE DEFICIENCY

Universal salt iodization is one of the most cost effective strategies in preventing Iodine deficiency disorders. The Quality and Standards Authority of Ethiopia, has set the iodine level to be 60-80 PPM as potassium iodate, after making allowance for losses of iodine during storage and distribution.



## 2.8.2. IRON

### Symptoms and Sign of the deficiency

#### Symptoms

Weakness

Fatigue

Dyspnea on exertion

Coldness and parasthesia of the hands and feet

Vague GI complaints such as

- Capricious appetite
- Flatulence
- Epigastric distress
- Belching
- Constipation or diarrhea

Glossitis (varying degree of papillary atrophy and soreness) which is

- More common over 40 years of age
- More in women than men

Angular stomatitis in 10-15%

Plummer Vinson syndrome in middle age women

#### Signs

Pallor

Pica (geophagia, clay eating and ice eating)

Finger nails and toe nails –lusterless, thin, brittle, flattened and spoon-shaped (koilonychias)

“Hemic” murmur, splenomegally, dependent edema, and lastly congestive heart failure.

## Diagnosis

Difficult to diagnose in mild cases

I. Clinical signs and symptoms

II. Biochemical Method

Hemoglobin determination or hematocrit

Peripheral morphology shows hypochromic microcytic

Serum Iron and serum ferritin decreases

Total Iron binding capacity increases usually greater than 84%

Bon marrow aspiration shows low iron storage in macrophages, decrease in number of side oblasts

III. Dietary

Dietary history

24 hours dietary recall

Food frequency questionnaire,

### Differential diagnosis

Thalassemia minor (target RBC)

Sideroblastosis (Perl's prussian blue stain)

Chronic illnesses



## TREATMENT

In treating iron deficiency anemia we should also consider administration of folic acid (see Table 1 below).

Table 1. Treatment protocol for nutritional anemia

<b>Age group</b>	<b>Dose</b>	<b>Duration of Treatment</b>
<b>&lt; 2 years</b>	25 mg iron + 100 - 400µg folic acid daily	3 months
<b>2 – 12 years</b>	60 mg iron +400µg folic acid daily	3 months
<b>Adolescents and adults including pregnant women</b>	120 mg of iron + 800µg folic acid Daily	3 months

Source: WHO, BASICS,, UNICEF. Nutrition Essentials: A guide for health managers

Oral or parenteral administration of iron in the form of gluconate, succinate, sulphate . Iron dextran should be administered parentally for patients with:

Ulcerative colitis, regional enteritis, intestinal shunts, colostomy or ileostomy

Malabsorption syndromes

Uncooperating patient or severe intolerance of oral dose

### Complications of parenteral administration

Anaphylactic shock particularly during i.V administration

Skin staining at the injection sites

### 2.8.3. SIGNS AND SYMPTOMS OF VITAMIN A DEFICIENCY DISORDER (VADD)

Night blindness

Lack of tear secretions (due to destruction of goblet cell)

Changes in eyes with eventual blindness if deficiency is severe and untreated

Susceptibility to respiratory infections

Dry, rough skin

Changes in mucous membranes

Weight loss

Poor bone growth

Weak tooth enamel

Diarrhea

Corneal Xerosis(Dryness)

Corneal Ulceration/ Keratomalacia

Bitot's spots

Conjunctival Xerosis

Corneal scar

Slow growth

It is characterized clinically by Xerosis (dryness) of the eyes and night blindness, which are classified by the WHO as follows:

**Night blindness (**

(Respiratory, gastro intestinal, genito-urinary tracts, as well as the conjunctivae are also affected).

**Corneal xerosis (X2A)** : is seen as a wrinkling of the cornea.

**A Corneal ulcer/ Keratomalacia (X3A)** is ulceration and softening of the cornea where



Serum level and  
Plasma Vitamin A and plasma carotene

### III. Dietary

Dietary history  
24 hours dietary recall  
Food frequency questionnaire,

#### **Treatment:**

Doses of oral vitamin A (as retinal palmitate in oil). One dose immediately, the second dose the next day, and the third dose 2 weeks later.

- 0-5 months 50, 000 IU per dose
- 6-11 months 100,000 IU per dose
- 12<sup>+</sup> months 200,000 IU per dose.

#### **Prevention:**

##### **Vitamin A supplementation**

- I. **Universal Supplementation:** A single dose of vitamin A every 6 months in population where vitamin A deficiency is a problem of public health importance (see criteria above).

Target groups: Children 6-59 months and lactating mothers within 6 weeks of delivery.

The supplementation could be done through the six critical contacts of women and children with the health service unit:

- Antenatal care
- Delivery
- Postnatal care and family planning
- Immunizations
- Well baby and
- Growth monitoring/ promotion visits and
- Sick child visit

Other outlets such as EPI+ programs, Community IMCI and clinical IMCI contacts can also be used for supplementing vitamin A.

**II. Disease targeted supplementation:** Give a single dose vitamin A to children under five years at the first contact with the health professional as follows:

A case of Protein energy malnutrition on the : 1<sup>st</sup>, 2<sup>nd</sup> and 14<sup>th</sup> days (Therapeutic dose)

A case of measles: 1<sup>st</sup>, 2<sup>nd</sup> and 14<sup>th</sup> days (Therapeutic dose)

Cases of acute respiratory illness and diarrheal diseases: a single dose on the 1<sup>st</sup> day of contact

**III. Food Diversification:** Give nutrition education on the importance of consuming diversified dietary sources of vitamin A such as green leafy vegetable (Kale, Cabbage) and Yellow fruits and vegetable (Mango, Papaya, Carrot, etc.)

C. Food Fortification: Different food vehicles like sugar can be fortified with vitamin A.

#### 2.8.4. VITAMIN D DEFICIENCY SYMPTOMS

Rickets (a childhood deficiency disease): malformations of joints or bones, late tooth development, weak muscles, listlessness, double malleolus, Rachitic Rosary, Bowed legs and knocked knees, delayed Fontanel closure, craniotabes

Osteomalacia (adult rickets): pain in ribs, lower spine, pelvis and legs, muscle weakness and spasm, brittle, easily broken bones.

#### Diagnosis

**I. Clinical:** signs and symptoms

#### **II. Biochemical**

Reduced levels of Vitamin D forms in blood.

Decreased serum phosphate, decreased calcium, increased alkaline phosphatase, urinary hydroxyproline, PTH levels.

III. Biophysical: Bone X-ray

IV. **Dietary Method:** Dietary history, 24 hour dietary recall method, food frequency questionnaire

### 2.8.5. FOLIC ACID DEFICIENCY



<b>Age group</b>	<b>Dose</b>	<b>Duration of Treatment</b>
<b>&lt; 2 years</b>	25 mg iron + 100 - 400µg folic acid daily	3 months
<b>2 – 12 years</b>	60 mg iron +400µg folic acid daily	3 months
<b>Adolescents and adults including pregnant women</b>	120 mg of iron + 800µg folic acid  Daily	3 months

Source: WHO, BASICS,, UNICEF. Nutrition Essentials: A guide for health managers

## 2.8.6. ZINC DEFICIENCY

### Moderate deficiency:

Impaired taste and smell (hypogeusia)

Growth retardation in children and in fetuses (results in low birth weight)

Delayed sexual maturation and impotence

Alopecia

Rashes

Multiple skin lesions (Acrodermatitis enteropathica)

Glossitis

Stomatitis

Impaired immunity

Abnormalities of labor including preterm labor or precipitated labor

Post partum hemorrhage due to uterine atony

Blepharitis

Paronychia

Sterility

Low sperm count and hypogonadism

Delayed wound healing

Night blindness and eye lesions including photophobia

Impaired appetite and food intake





Growth monitoring and well baby clinic visits

Sick baby clinic visits

During all these contacts **Essential Nutrition Actions (ENA)** that need to be promoted are the following:

- I. Exclusive breast-feeding up to 6 months of age
- II. Complementary feeding at the age of 6 months with continued breast feeding up to 2 years and beyond
- III. Continued feeding of the sick child
- IV. Maternal nutrition
- V. Vitamin A supplementation for children 6-59 months and for lactating mothers within 8 weeks after delivery.
- VI. Iron and folic acid supplementation to the mother during pregnancy and in the postnatal period
- VII. Ensure the consumption of iodized salt by the family

### **2.12 Learning activities (case study) continued**

Refer to case studies 1,2 and 3 in the core module and discuss on the following questions in the class. The instructor can assist you.

1. *What pertinent history do you ask parents of children in cases 1.2 and 3?*
2. *What pertinent physical signs would you look for?*
3. *What laboratory investigations would you order in order to substantiate your diagnosis?*
4. *What other assessments do you carry out in order to determine the type of malnutrition?*
5. *What is your diagnosis from the story?*
6. *What other causes do you consider for the differential diagnosis?*



# UNIT ONE

## SATELLITE MODULE FOR PUBLIC HEALTH NURSES

### Introduction

Adequate nutrition is the intake and utilization of enough energy and nutrients, together with disease control, to maintain well-being, health and productivity. “Malnutrition” includes generalized malnutrition (which manifests itself as stunting, underweight and wasting in individuals) and deficiencies of micronutrients such as vitamin A, Iron, Zinc, and Iodine. The magnitude of the problem relating with micronutrients in Ethiopia are so intense and majority of children, adults, pregnant and lactating women are affected. Therefore, health workers particularly the public health nurses are critical to the design and implementation of nutrition program. Collaboration with the other professionals and implementing the essential nutrition action can reduce infant and child mortality; improve physical and mental growth, development, and productivity.

#### 1.1. Purpose and Use of the Satellite Module

The purpose of this satellite module is to equip learners (trainees) with knowledge and skills required to identify and manage effectively cases of micronutrient deficiency. The public health nurses can use this satellite module in their-services or in-services training programs.

#### 1.2. Direction for using the Satellite Module

For a better understanding of this module, the public health nurses are advised to refer to the (prtfes 28oa2T

# UNIT TWO

## SATELLITE MODULE FOR PUBLIC HEALTH NURSES

### 2.1 Pre and Post test

See the core module unit 2

### 2.2 Significance and Brief Description of the Problem

See the core module unit 2.

### 2.3 Learning Objectives

To be able to prioritize the key nutrition behaviors which met the health and nutrition needs of vulnerable groups of population in the communities and integrate these in to ongoing intervention programs in health facilities and communities at large.

### 2.4 Case Study: Learning Activities.

Read cases 1, 2 and 3 so that you will be able to discuss questions in section 2.12 of this module.

### 2.5 Micronutrient Deficiencies:

Micronutrient deficiency refers to the deficiency (decrease in the level) of micronutrients that are essential for the vital function including growth, development, immunity, fertility

and other lively processes. Although there are various micronutrients essential to our body, in this regard, this module focuses mainly on few micronutrients that are of public health importance in the Ethiopian context due to the tolls of morbidity, disability and mortality they are causing in the population. Therefore, this module focuses mainly on the following micronutrients: Vitamin A deficiency, Iron Deficiency anemia, Iodine Deficiency disorder and vitamin D deficiency (rickets),

### **2.5.1. VITAMIN A DEFICIENCY**

Vitamin A has the distinction of being the first fat-soluble vitamin to be recognized. It is certainly one of the most versatile, with parts to play in such diverse functions as vision, maintenance of body lining and skin, bone growth, anti oxidant activity and reproduction. A deficiency may be as many as hundred thousand a year world wide, placing a heavy burden on society.

Until recently vitamin A deficiency was thought to be a problem only for the eye health and vision. It is now clear that it is also essential for the proper functioning of the immune system of the children, which occurs, long before damage to the eye is apparent.

It is also worth to mention that the vitamin A status of the infant is closely correlated with that of the mother. Vitamin A deficient mother transfers less vitamin A to their fetuses. After birth the concentration of vitamin A in the breast milk depends on the vitamin A status of lactating woman. Breast milk depends on the vitamin A status of the mother.

## **Vitamin A:**

Important component for seeing in dim light.

Has to come from food or supplements

Is stored in the liver

Builds the epithelial cells

Is important for immune system.

## **Who is at the greatest risk?**

Children 6 to 59 months

Women during pregnancy and lactation

## **Objectives:**

Eliminate vitamin A deficiency

Decrease child morbidity and mortality.

## **Solution to vitamin A deficiency**

### **Strategies**

Breast milk

Food diversification

Supplementations

Food fortification

## **1. Breast feeding**

Breast milk is a hygienic source of energy, essential nutrients, water, immune factors, and many other components that are beneficial for infant and young children. Breast milk protects against vitamin A deficiency. Breast milk is the ideal food for the children in the first six months of age. This is because:

Breast milk is rich in vitamin A


Exclusive breastfeeding reduces infection and vitamin A losses.

Postpartum vitamin A supplementation of lactating women will raise breast milk vitamin A content.

Promotion of exclusive breastfeeding is a strategy for preventing Vitamin A deficiency.

Vitamin A intervention increases the benefits of breast feeding promotion for maternal and child health and survival.

The Public Health Nurses should always utilize the following important key message at each contact with the mother in order to promote breast feeding.



## Treatment supplementation for different diseases

Pneumonia, severe malnutrition, severe anemia and persistent diarrhea Capsule on day 1.

Measles: capsule on day 1, day 2, day 30.

Treatment of eye problems due to Vitamin A deficiency: capsule on Day 1, Day 2. Day 30.

**4. Food Fortification:** Adding vitamin A to foods that are commonly consumed by the high risk groups, such as: Vitamin A enriched sugar, Iron enriched flour, fish liver oil ... etc..

## Key contact points for control of Vitamin A deficiency.

- Pregnancy
- Delivery
- Postnatal and family planning
- Well child/ GM/P
- Immunization
- Sick child

## Pregnancy (antenatal contact)

Increased food and vitamin A intake. In the areas where vitamin A deficiency is common and vitamin A rich foods are scarce, low dose of vitamin A supplements (< 10 000 IU/ day or 25 000 IU/week) or multiple micronutrient supplements with appropriate levels of vitamin A may be recommended.)

Counsel on early initiation of breast feeding, feeding of colostrum, and frequent exclusive breast feeding

Assess nutritional status and encourage increased energy intake, a varied diet, including vitamin A-rich food, and reduced workloads during pregnancy

Discuss family planning options that protect lactation, including Lactational Amenorrhea Method (LAM)

Educate on parasite prevention and treat parasite infection.



## **Immediate postpartum Contact**

Facilitate early initiation of breast feeding, counsel on frequent and exclusive breast feeding, and help establish good breast feeding skills (Proper positioning and attachment).

Administer a single high dose vitamin A supplement to mother immediately after delivery.

Assess nutritional status and counsel mother on increasing energy intake, consuming a varied vitamin A rich diet, and reducing work load through out lactation.

Discuss family planning options that protect lactation, including LAM.

## **Postnatal contact**

- Vitamin A deficiency will result in low vitamin A concentration in breast milk.

In areas where vitamin A deficiency is common, give a single high dose [200,000 IU] vitamin A capsule as soon after delivery as possible, but no later than eight weeks postpartum. This will help to build up vitamin A stores, improve the vitamin A content

### Well baby contact (Growth monitoring and immunizations)

Provide nutritional assessment and counseling, reinforcing the mother's need for increased energy intake, a varied vitamin A diet, and reduced work load.

Counsel on exclusive breast feeding, timing for the introduction of appropriate complementary food, the importance of vitamin A rich foods in the child's diet, and practical ways of enriching the child's diet using locally available foods.

Check and complete vitamin A supplementation protocol for older infants and young children.

Check and complete mother vitamin A supplementations (If within 8 weeks postpartum)

Discuss family planning options that protect lactation, including LAM, IUCD...etc.

Administer 50,000 IU Vitamin A supplement to non breast fed infant under six months at first contact

### The sick baby clinic/ Sick child:

Assess and counsel on breast feeding and adequate and complimentary feeding during and after illness.

Check and complete vitamin A supplementation protocol: Follow national policies therapeutic dosing with high dose of vitamin A capsule for measles, xerophthalmia, chronic diarrhea and severe malnutrition.

### **Action to support breast-feeding and improve vitamin A status:**

Use local communication channels (mass media, schools, religious institutions, community events, market place, community based distributions systems etc.) to disseminate information on optimal infant feeding practices, maternal nutrition, and vitamin A.

Increases availability of vitamin A-rich foods through schools and community gardens, fruit trees and animal husbandry (raising hens, chickens, and other small animals).

## 2. Household level

The public health nurse should advise the mother at home setting level that sun dried seasonal fruits and vegetables or air dried out of direct sunlight would provide a source of vitamin A through out the year.

Recommend the mother in a way that she may be capable of serving vitamin A-rich foods with a little fat (oil, butter, peanuts) to increases utilization by the body.

Recommend communities to plant gardens with a variety of vitamin A-rich vegetables such as Mango, Sweet potatoes, Pumpkin, Carrot

### Basic Facts: Consequences of Vitamin A deficiency in Pregnancy

#### Increases the risk of:

- Night blindness
- Maternal mortality
- Miscarriage
- Still birth
- Low birth weight
- Reduced transfer of Vitamin A to fetus

### 2.5.2 Iron Deficiency Anemia

Anemia is low level hemoglobin in the blood, as evidenced by reduced quality or quantity of red blood cells that affects oxygen carrying capacity of blood. It has massive consequences for maternal and child health, child development and productivity. Iron is essential to periods of growth, such as infancy, adolescence, pregnancy and lactation.



Premature birth  
Intrauterine growth retardation  
Low birth weight  
Increased risk of maternal mortality  
Reduced ability to survive bleeding during and after birth  
Decreased productivity due to fatigue and tachypnea.

### **Consequences of Anemia in Children:**

Increased infant mortality(due to low birth weight)  
Learning deficits (Iron Deficiency anemia lowers IQ by 10 points)  
Fatigue and poor growth  
Iron deficiency affects iodine up take thus increasing risk of Iodine Deficiency Disorders

### **Critical Contact Points**

Essential nutrition action needs to be taken by all health and extension workers.

Those critical contact points are:

During pregnancy  
During delivery  
During postnatal period  
Well baby clinic  
Immunization  
Sick child clinic

#### **1. Antenatal contact**

Give Iron /folic acid supplementation  
De-worm pregnant women  
Give anti malarial to pregnant women  
Promote mosquitoes-nets for mothers and baby  
Promote improved diet for women  
Prepare for optimal breast feeding once the baby is born

Promote consumption of iodized salt during antenatal Visits.

## **2. Delivery Contact**

Put the baby to breast immediately

Give Vitamin A to mother

Continue` Iron/folic acid for the mother

Promote improved diet for mother during lactation

### **Post-natal and/ or family planning contact**

Promote exclusive breast feeding for 0-6 months

Check appropriate position and attachment during breastfeeding

Promote improved diet for mother

Continue iron/folic acid for mother

Promote mosquito-nets for mothers and baby

Promote consumption of iodized salt and FP, STI prevention and child's vaccination.

## **3. Immunization Contact**

Complete Vitamin A for the baby

De-worm older infants

Assess and treat infants anemia

Promote optimal breast feeding exclusive for 0-6 months

Promote adequate complementary feeding

Promote use of mosquito-net mother and child` Vaccinations (child and women) and FP, STI and timely referral.

## **4. Well child visit and GM/P Contact**

Monitor infant/child's growth

Assess and counsel on feeding BF and CF (breast feeding and complementary feeding)

Assess and treat infant anemia

- Promote consumption of iodized salt
- Complete vitamin A
- De-worm for parasites
- Check and complete vaccinations.

## 5. Sick child visit (IMCI) contact

- Monitor infant /child growth
- Counsel on feeding breast feeding and complementary feeding
- Assess and treat anemia
- De-worm
- Complete vitamin A and vaccinations
- Assess and treat as IMCI

## 6. Contact in other sectors

### 6.1. In school Programs

- Assess and treat child and adolescent anemia
- Promote consumption of iodized `salt
- Complete vitamin A
- De-worm
- School lunch if possible
- Promotion of better nutrition

Assess and treat parasites

### 6.2 In out reach sites

- Monitor infant /child growth
- Promote breast feeding
- De-worm for parasites

### **6.3 In existing community based organization meetings.**

Promote adequate breast feeding and complementary feeding

Promote control of anemia

Promote consumption of iron rich foods

Promote consumption of Vitamin A rich food

Promote consumption of iodized salt

Promote vitamin A supplementations

Promote increased food intake during pregnancy and lactation

National immunization days for polio and measles can be used to give Vitamin A.

### **6.4 In Micro-credit groups targeting women:**

Promote adequate breast feeding and complementary feeding

Promote control of anemia

Promote consumption of iron rich foods

Promote consumption of Vitamin A rich food

Promote consumption of iodized salt

Promote vitamin A supplementations

Promote increased food intake during pregnancy and lactation

### **6.5 In Emergency situation: In addition to therapeutic feedings and**



Promote environmental sanitation and clean water.

### **2.5.3 Iodine Deficiency Disorders (IDD)**

Proper nutrition is essential for mental and physical development and school performance. Evidence from different countries shows that malnutrition considerably reduces children's learning ability, school performance and retention rates. Iodine deficiency lowers the ability of children to think and become creative and productive adults. Iodine is also necessary for the normal development of the brain of fetus during pregnancy. Pregnant women living in iodine deficiency regions are more likely to give birth to mentally retarded children.

IDD remains a significant public health problem in many countries. According the 1990 nutritional survey in Ethiopia, the total rate of goiter, the most visible sign of iodine deficiency is 22 %. Globally, 30 % of the world's population is affected by IDD and more than 150 million people are affected in Africa alone. In Ethiopia, one out of every 1000Tw 2arrably

In school age children

Women in the reproductive age

Men and females of all ages

Young children and growing fetus are particularly vulnerable to the damage caused by IDD.

Infants who are not exclusively breast fed

Lactating women

### **Strategies used to control IDD**

1. Distribute iodized oil capsule
2. Increase access and consumption by all families of iodized salt

Distribution of Iodized oil capsule

The public health nurse should distribute to all potential clients if available.

For High endemic areas only the following regime should be utilized

1. capsule for pregnant women and children less than 5 years
2. Capsule for women and children for 5 to 14 years of age.

### **Key contact points for control of IDD**

Pregnancy

Delivery

Postnatal and family planning

Immunization

Well child/ GM/P

Sick child Clinic

-(Refer the essential activities on the key contact points for Vitamin A and Iron deficiency as applied to IDD)

-The public nurse should know the following important element of a successful IDD program.

### **Elements of a successful IDD program**

Supplement supply  
Delivery system  
Demand and compliance  
Monitoring and evaluation

#### **2.5.4 Vitamin 'D' Deficiency (Rickets):**

Rickets is a disease caused due to Vitamin D deficiency. It is characterized by weakness and deformity of bones. Deficiency states occur due to lack of calcium ion in the body which internally is mainly due to defective absorption because of vitamin D deficiency.

The two most important vitamins 'D's are vitamin D-2 and D-3. Vitamin D-2 (ergo-calciferol) exists in fungi and yeast and in the form generally added to milk, food and vitamin supplements. It is used to combat hypocalcaemia (low blood calcium). Vitamin D-3 (cholecalciferol), is found in fish oils, egg yolks and milk. The body produces vitamin D-3 in the presence of ultraviolet light.

Unhealthy child care practices such as non exposure of infants to sunshine and complete covering while outdoors for fear of evil eyes contribute to the occurrence of rickets in Ethiopian children.

Since sunshine is adequately available in Africa, rickets is uncommon. But in countries like Ethiopia, traditional and cultural practice of child-care has contributed to the occurrence of rickets. Rickets is more commonly associated with protein energy malnutrition and improper weaning practice.

#### **Functions of Vitamin D**

Regulates growth, hardening and repairing of bone by controlling absorption of calcium and phosphorus from small intestine.

Prevent rickets

Prevent hypocalcaemia (low blood calcium)

Prevent post-operative muscle contraction



Sardines

Sunlight plus subcutaneous fat

Vitamin D fortified milk.

### **How to take available capsule or tablets**

Take with or immediately after food to increase absorption

Take oral solutions with meals or 1-1: 30 hours after meal.

High dose of vitamin D have been generally discouraged as it has side-effects including: calcification of soft tissue, kidney failure which is manifested by high blood pressure, irregular heart beat, nausea, weight loss, seizure, mental and physical growth retardation, kidney damage ...etc

## **2.6 Practical Nursing for Improving Nutrition**

Guidelines for appropriate complementary feeding of infants and young children 6- 24 months.

***Appropriate complementary feeding promotes growth and prevents stunting among children 6-24 months.***

The period of complementary feeding is when others foods or liquids are provided along with breast milk. Rates of malnutrition's usually peak at this time with consequences of persist throughout life. Stunting is seldom reversed in later childhood and adolescents. Inadequate feeding of female children also affects nutrients stores, subsequent reproductive health, and the risk of maternal mortality.

***Appropriate complementary feeding invol***



Feed energy-dense combination of foods to 6-11 months olds.

Introduce “finger foods”( snacks that can be eaten by children alone) beginning around 8 months of age in the mean time, try to avoid risk of chocking

Make the transition to the family diet at about 12 months of age.

### ***Diversify the diet to improve quality and micronutrient intake.***

Feed Vitamin A-rich fruits and vegetables daily.

Feed meat, poultry, or fish daily or as often as possible, if feasible and acceptable.

Use fortified foods, such as iodized salt, Vitamin A-enriched sugar, iron enriched sugar, Iron enriched flour or other staples, when available.

### ***Practice active feeding***

Feed infants directly and assist older children when they fed themselves.

Offer favorite foods and encourage children to eat when they lose interest or have depressed appetites.

If children refuse many foods, experiment with different food combination, tastes, texture, and methods for encouragement

Talk to children while feeding.

Feed slowly and patiently and minimize distractions during meals.

Do not force children to eat.

### ***Practice frequent and active feeding during and after illness.***

During illness, increases fluid intake by more frequent breastfeeding, and patiently encourage children to eat favorite foods.

After illness, breastfeed and give foods more often than usual, and encourage children to eat more food at each sitting.

### ***Practice good hygiene and proper food handling***

Wash caregivers' and children's hands with soap before and after food preparation and eating.

Serve foods immediately after preparation

Use clean utensil to prepare and serve food.

Serve children using clean cups and bowls, and never use feeding bottles.

i. Responsibility of the public health nurse to





This involves teaching all sections of the community, especially fathers and mothers, to make the best use of the foods available (including breast feeding), to make use of available primary health care services, and to grow local foods in their own garden.

**There are seven rules, which, if kept, can largely improve nutritional status in the community.**

1. Identify the local sources of food stuffs.
2. Recognize the causes of improper feeding in the community.
3. Explain the effects of improper feeding on different age and sex groups.
4. Teach nutritional food values of local food stuffs.
5. Demonstrate how to cook balanced meals using locally available foods.
6. Teach personal and food hygiene at home.
7. Evaluate what the community members have learned about improved nutrition in order to prevent micronutrient deficiency
8. Make the recommended meals to prevent effects of improper

### Signs of properly attached infant:

- a. Mother brings infant toward her breast, not the breast toward her infant
- b. Infant's mouth is open wide.
- c. Infant's lower lip turns outwards
- d. Infant's chin touches mother's breast
- e. Mother entire nipple and a good portion of the areola ( dark skin around the nipple) are in infant's mouth.

### Mother initiates breastfeeding within 1 hour of birth.

Takes advantages of the newborn's intense suckling reflex and alertness

Stimulates breast milk production

Protects infants from diseases by providing the thick yellowish milk (colostrums),

Helps to expel placenta more rapidly and reduce blood loss.

Helps expel meconium the infant's first stool.

Keeps newborn warm through skin-to skin contact

Fosters bonding between mother and infant

Mother breast feeds frequently, day and night.

Mother offers second breast after infant empties the first.

Mother continues breastfeeding when either she or the infant is sick.

Mother who will be away from her infant for an extended period expresses her breast milk. Care givers feeds expressed breast milk from a cup.

Mother expresses breast milk following these steps.

washes hands

- Prepares a clean container
- Gently massages breasts in a circular motion
- Positions her thumb on the upper edge of the areola and the first two fingers on the underside of the breast behind the areola.
- Pushes straight into the chest wall.
- Avoids spreading the fingers apart.
- For large breasts, first lifts and then pushes in to the chest wall

- Rolls thumb and fingers forward as if making thumb and fingerprints.
- Repeats rhythmically: position, roll; position; push, roll
- Rotates the thumb and finger positions.

#### Learning Activities (Case Study) Continued

Refer to the story of cases 1,2 and 3 in the core module and discuss on the following questions among the team.

1. What type of major health problem did the health center team identify in that particular community?
2. What fundamental intervention programmes need to be planned by the health workers in general? ( Consider all the given case scenario)
3. Who should be involved in identifying and prioritizing the health problems for better intervention and good outcome?
4. What is expected from the health workers as far as the given case scenario is concerned?
5. What type of feeding behavior and practices would bring change and improve the health of the community?
6. What basic things were thought by the nurse in order help children to grow healthier and to prevent micronutrient deficiency?
7. What will happen to children if they do not get the necessary nutrients?
8. What will be the role of the public health nurse in the promotion of health and prevention of diseases in the community?

# UNIT ONE

## SATELLITE MODULE FOR LABORATORY TECHNICIANS

### Introduction

#### 1.1 Purpose of the module

This module helps laboratory technicians to participate in the team in the diagnosis and management of micronutrient deficiencies, with a particular emphasis on the laboratory investigations of selected micronutrients such as iron, iodine, zinc, folate, vitamins A and D. Additionally useful laboratory tests associated with these micronutrient deficiencies are discussed.

#### 1.2 Direction for using the satellite module

For a better understanding of this module the laboratory technicians are advised to follow the following directions.

Do the pretest in your profession in unit 2, section 2.1.2.3 of the core module

Read the core module thoroughly

Use listed references and suggested reading materials to substantiate your understanding of the problem

Evaluate yourself by doing the post test and referring to the keys given in unit 7 section 7.1.2.3

# UNIT TWO

## SATELLITE MODULE

### 2.1. Pretest

*Refer to the pre and post test in the core module unit 2 section 2.1.2.3*

### 2.2. Significance and brief description of the problem

*See the core module unit 2 section 2.2.*

### 2.3. Learning Objectives

After completion of this module students will able to:

Describe how to collect, handle and label blood specimens

List the different specimens that are used for the diagnosis of micronutrient deficiencies.

Describe the concept of laboratory diagnosis of micronutrient deficiencies.

List biochemical tests used in the diagnosis of iron deficiency anemias

Describe the micro method of packed cell volume determination.

Describe and demonstrate how to prepare and stain thin blood film for red blood cell morphology

Demonstrate how to assess stained thin blood films including elements of the blood films other than red cell morphology (e.g. hemoparasites)

Classify anemia based on red blood cell morphology and measured hemoglobin

Indicate the biochemical profiles of microcytic hypochromic anemia due to iron deficiency

## **2.4 Learning activities: Case study**

## **2.5 Definition**

*Refer to the core module unit 2 sections 2.5.*

## **2.6 Epidemiology**

*Refer to the core module unit 2 sections 2.6.*

## **2.7 Cause, Etiology and pathogenesis**

*Refer to the core module unit 2 sections 2.7.*

## **2.8 Clinical features**

*Refer to the core module unit 2 sections 2.8.*

## **2.9. Diagnosis**

### **2.9.1 Introduction**

Medical laboratory professionals are mainly

T<sub>3</sub> in dried blood spots exposed to









### 2.9.3. Estimation of hemoglobin by the Acid Hematin Method of Sahli-Hellige

**Principle: -**

*Hemoglobin in a sample of blood is converted to a brown colored acid hematin by treatment with 0.1 N HCl and after allowing the diluted sample to stand for 5 minute to ensure complete conversion to acid hematin it is diluted with distilled water or 0.1 N HCl until its color match as with the color of an artificial standard (tinted glass).*

Depending on the type of hemoglobinometer, this gives the hemoglobin concentration either in g/dl or as a percentage of normal where 100% is equal to 14.6 g/dl. To convert the percentage to g/l, multiply by 1.46 or to g/dl by 0.146. Other standardized haemoglobin estimation methods could be used whenever available. Though, expensive the haemo-cue method is more reliable and dependable than the Sahli's method. As some other colorimetric methods it is not technically demanding, standardization is simple and operation with out direct electric supply option is offered.

**Normal range of hemoglobin at different age groups**

Hemoglobin in mg/dl	
Children at birth	13.6-19.6
Children at 1 year	11.3-13.0
Children, 10-12 years	11.5-14.8
Women	11.5-16.5
Men	13.0-18.0

**Source: Harriet Lane handbook**

## 2.9.4. PREPARATION, STAINING AND EXAMINATION OF PERIPHERAL BLOOD FILM

### 2.9.4.1 Preparation of Thin Blood Film

Examination of the blood film is an important part of the hematological evaluation and the validity or reliability of the information obtained from blood film evaluation, the differential leukocyte count in particular depends heavily on well-made and well-stained films. If not made from skin puncture, films should be prepared within 1 hour of blood collection into EDTA. Adequate mixing is necessary prior to film preparation if the blood has been standing for any appreciable period of time. A thin blood films can be prepared on glass slides or cover glasses. The latter has the single most important advantage of more even distribution of leucocytes. Preparation of blood films on glass slides has the following advantages:

- Slides are not easily broken

- Slides are easier to label

- When large numbers of films are to be dealt with, slides will be found much easier to handle.

**Generally it is advisable to make a smear that covers about 2/3 or 3/4 the size of the slid. Cells** should evenly spread in monolayer, especially at the examination area near the tail of the smear. Preferably blood slides should be labeled before they are made. The commonest practice is, once the smear is dry, it should be labeled appropriately with the name of the patient and date or a reference number on the head of the film using a lead pencil or graphite. If these are not available, labeling can be made by scratching the above information on dried smear with the edge of a slide. A paper label should be affixed to the slide after staining.

### 2.9.4.2 Staining of thin blood film with Romanowsky dyes

Modern Romanowsky stains in common use (e.g., Wright's, Leishman's) containing an acidic component (eosin B) and a basic component (methylene blue) are commonly used. Also panoptic stains like MayGrunwalds- Giemsa could be used to stain blood films for differential leukocyte count and red cell mo





### **2.9.5.1 Methods of counting**

Various systems of performing the differential count have been advocated. The problem is to overcome the differences in distribution of the various classes of white cells, which are probably always present to a small extent even in well-made films.

The lateral strip (“crenellation”) pattern of differential counting is the most routinely used pattern and in this method the field of view is moved from side to side across the width of the slide in the counting area just behind the feathered edge where the cells are separated from one another and are free from artifacts. Multiple manual registers or electronic counters are used for the count.

N.B. The following elements of the blood film must be observed while performing the differential count.

Erythrocytes: size, shape, degree of hemoglobinization, presence of inclusion bodies

The presence of atypical lymphocytes

Haemoparasites: malaria, borrelia, babesia, microfilariae, trypanosoma, etc.

### **2.9.5.2 Reporting the differential leukocyte count**

The differential leukocyte count could be expressed as the percentage of each type of cell or it could be related to the total leukocyte count and the results reported in absolute numbers.

### 2.9.5.3 Normal differential ranges:

	1-4 years	10 years	Adults
Neutrophils	36-48%	45-55%	55-65%



of red cells to whole blood in liters or in percentage. By microchaematocrit it is only possible to measure results to two decimal places.

The major role of the primary health care laboratory professional in the diagnosis of micronutrient deficiency is to perform tests used in presumptive diagnosis. Additionally for specific diagnosis using biochemical tests involved in proper collection, preparation, storage and transportation of the appropriate samples to the central or specialized laboratories. The type of specimen, storage and transportation requirement may vary according to the test protocol used, hence following the instructions is mandatory. However, limited hematological investigations such as mentioned above could be performed to aid in the diagnosis of iron deficiency anemia.

## **2.10 Case management**

*Refer to the core module unit 2 sections 2.10.*

## **2.11. Prevention and intervention**

*Refer to the core module unit 2 sections 2.11.*

## **2.12. Learning activities (case study) continued**

Refer to story of health professionals in the core module and discuss on the following questions in the class. The instructor can assist you.

*How is blood specimen collected, stained and examined for blood morphology examination?*

*What could be the etiology of micronutrient deficiencies?*

*What laboratory investigations could be done at the health station or health center level?*

*What should be reported in the laboratory request form in the determination of hematological tests for the assessment of iron and folate deficiency?*

*What is the prime role of laboratory professionals at primary health care level in the diagnosis and management of micronutrient deficiencies?*



### **2.13 Roles and Task Analysis**

*Refer to the core module unit 4.*

### **2.14 Glossary and abbreviations**


*Refer to the core module unit 5.*

### **2.15 Bibliography**

*Refer to the core module unit 6.*

### **2.16. Annexes**

*Refer to the core module unit 7.*



# UNIT ONE

## SATELLITE MODULE FOR SANITARIAN

### Introduction

The role of the sanitarian in the prevention of micronutrient deficiencies is mostly in awareness creation, environmental sanitation improvement and behavioral changes in nutritional improvement and hygiene practices.

### 1.1 Purpose and use of the module

The main purpose of this module is to equip the sanitarians with adequate knowledge and skills for the prevention of micronutrient deficiencies together with other team members.

### 1.2 Directions for using the module

For a better understanding of this module, the sanitarian is advised to follow the following directions

- Do the pre test pertinent to your field in unit 2 of the core module (The sanitarian should also read the core module thoroughly at first and when referred in this module)

- The sanitarian could be successful in using this module if he works with other team members and intersectorally with other development workers (agriculture extension workers, development workers, home economists, etc)

- Evaluate your self by doing posttests and comparing your score with the pre tests.

## UNIT TWO

### SATELLITE MODULE

#### 2.1 Pre test and posttest:

Please refer to the core module

#### 2.2 significance and brief description of the problem:

Please refer to the core module

#### 2.3 Learning objectives

At the end of reading this module the sanitarians will be able to:

Describe the prevention methods of micro nutrient deficiencies

Identify appropriate methods and the primary targets for nutrition and health education program in the prevention of micro nutrient deficiencies

Describe why personal hygiene, environmental sanitation and nutrition education prevent those risk factors which are associated with micro nutrient deficiencies

Describe the whole mechanism of different factors that are associated with the problem of micro nutrient deficiencies

#### 2.4 Learning activities:

R0.0re mp R0.0r4tm mu7Dca< 4Be.0021 Tw -o >BDC

high cost of foods rich in iron (refer to case study 1). Dietary diversification can be addressed by:

- i. *Economic approach* (promotion of agriculture). This approach aims at improving the incomes of the target community as a solution to their nutritional problems.
  - ii. *Income generation scheme*: Income generation schemes enable the target community to buy food. It includes organizing the community and using their potentials in the running of the project. The target communities according to their needs or resources available could identify the project. The project may be weaving, bee keeping, pottery, etc.
2. *Fortification*: The fortification of staple foods such as wheat or other grains is likely to increase iron intake for those population that have access to them. However, infants and children who have a limited capacity to eat large quantities of fortified food are not likely to benefit significantly from this strategy. Targeted fortification as is practiced by the developed world is an excellent way to increase the intake of iron in children. Nevertheless, fortified commercially prepared infant foods are relatively expensive and may not be affordable for many families with children at highest risk of iron deficiency.
3. *Supplementation*: The final approach is through supplementation of individuals or communities at risk. For infants and children, this approach is the most likely to succeed. Supplementation may be the best way to reach infants and young children in families that can't afford expensive iron containing or fortified foods. Challenges from supplementation, however, includes compliance and adequate distribution of the supplement.
4. *Environmental sanitation*: Many studies have shown that iron deficiency anemia is associated with malaria and other parasitic infections. Therefore, to prevent these problems the following are major interventions that has to be conducted by the sanitarian together with other team members in the community.



existing behavior. Each culture lives in a unique environment requiring evaluation to understand the interaction contributi





## What needs to be done?

The technology for adding potassium iodate to salt during processing and refinement is one of the lowest-cost health and nutrition interventions available.

In summary, an array of nutrition interventions is available to address nutrition problems in different settings. Based on experiences in the past two decades, the most cost-effective, widely applicable, and manageable nutrition interventions for primary health care services consist of promoting, protecting, and supporting-

- Exclusive breastfeeding for at least four, and if possible, for six months;
- Appropriate complementary feeding and continued breastfeeding to two years
- Adequate nutritional care of sick and malnourished children;
- Adequate vitamin A intake;
- Adequate iron intake;
- Adequate iodine intake

To assess the existence of micronutrient deficiency in a given community the sanitarian should assess the nutritional situation of the community using different indicators. Use the following table as a guideline in collecting data.

Examples of household survey questions for priority nutrition interventions (WHO, 1999).

Priority behavior	Indicator	Questions
Exclusive breast feeding	% of infants 0-6 months of age who are exclusively breastfed	What did the child eat yesterday? (Note if anything other than breast milk was given except vitamin drops.
Complementary feeding and continued breastfeeding to two years	% of infants 6-9 months of age given breast milk and semi-solid complementary foods. % of children 6-23 months who are actively encouraged to eat. % of infants 12-18 months of age given semi-solid or solid	What did the child eat yesterday? Was this food liquid, semi-solid, or solid? (Record the number of times the child was fed). What ingredients were in the child's food? (Use a checklist of locally available foods rich in vitamin A and C,



	<p>complementary foods at least 4 times/day, in addition to breast milk.</p> <p>% of children 20-23 months of age who are breast feeding</p>	<p>local sources of animal products, and energy-dense foods.)</p> <p>Did you actively encourage the child to eat yesterday? If yes, what did you do? (Correct answers: Give foods liked by the child, sat with the child, and others.)</p>
--	--	--

Care of sick children

% of children 0-23 months of age who were sick in the past 2 weeks, and increased breast feeding.

% of children 0-23 months of age, who were sick in the past 2 weeks, & didn't reduce feeding other foods.

% of children 6-23 months of age sick in the past 2 weeks and were encouraged to eat.

Was the child ill in the past 2 weeks?

If yes, did the child breastfeed more, less or the same?

If yes, did the child eat more, less or the same amount of other foods?

If yes, did you actively encourage the child to eat during or after the illness? (Correct answers: Gave special foods



public education can solve vitamin A deficiency problem that is common throughout the nation.

*Fortification:* For people who can afford to pay adding vitamin A to foods that are commonly consumed by the high-risk groups.

*Supplementation:* In vitamin A deficiency risk areas, giving age-appropriate doses of oral supplements of vitamin A to children and to women (within the first 6-8 weeks) after delivery is an option if available.

*Preventing illnesses:* Preventing illness and treating them early also protects against reduced vitamin A stores and contributes to reducing VADD.

Preventing illnesses also protects against VADD.

# UNIT ONE

## SATELLITE MODULE FOR HEALTH EXTENSION WORKERS (HEW)

### Introductions

#### 1.1. Purpose & Use of the Module

Materialization of the Community based management of micronutrient deficiencies is made possible through training of HEW that are well equipped

-15knosc ski-18jmco

# UNIT TWO

## SATELLITE MODULE FOR PRIMARY HEALTH WORKERS (HEW)/COMMUNITY HEALTH WORKERS

### (HEW)

#### 2.1. Pre and posttest

See the pre and post test for primary health workers (HEW)/Community health workers (HEW) in the core module section 2.1.2.5

#### 2.2. Significance and brief descriptions of the problem

*The user of this module for training HEWs is highly advised to refer to the core module sections 2.2.*



- **Learning Objectives**

- At the end of completing this module the HEWs will be able to:

Define and identify types of micronutrient deficiencies

Identify symptoms and signs of common micronutrient deficiencies

Demonstrate preparation of nutritious food that contain the essential micronutrients to the mothers and care givers

Refer children with micronutrient deficiencies to the to the appropriate health institution

Give health education on the preventive methods of micronutrient deficiencies and importance of child nutrition for proper growth and development

Advise mothers/care givers on the importance of exclusive feeding during the first 6 months and complementary feeding with breast milk there after

Educate mothers/care givers/or other members of the family about the importance of horticulture and backyard gardening, immunization, importance of continued feeding during diarrhea



#### **2.4. Learning activities (case study)**

Read the different stories in the core module for the class or make them read it thoroughly so that they will be able discuss the questions related to the story in unit 2, section 2.12

## 2.5. Definition

**Micronutrient** mal nutrition is the manifestation of deficient dietary intake of micronutrients that provide the body with the function of growth, development, immunity, fertility, and other essential bodily activities that are dependent on the supply of the particular nutrient.

## 2.6. Epidemiology

Micronutrient deficiency is called a hidden hunger because it is unnoticeable and has far-reaching consequences. Besides there are simple solutions to it. In Ethiopia, about 40% of all mothers and 50% of pregnant mothers are anemic. Vitamin A deficiency affects about 27% of the population. Total goiter rate is 22%. As the diet of over 85% of the society is based on cereal foods, there the deficiency of other micronutrients like zinc and iron.

## 2.7. Causes

Different factors contribute to the occurrence







- History of immunization
- Birth interval in the family
- Child care practices

### **Physical Examination**

- Vital signs –Pulse rate, Respiratory rate, Weight and height
- Check for specific signs of micronutrient deficiency
- Check for the underlying causes of the anemia (infection, malabsorption, hemorrhage, etc.)

### **2.10. Case management**

Upon regular **growth monitoring** care givers of those children with micronutrient deficiencies should be educated to improve their child feeding practices by preparing a balanced diet from locally available foodstuffs. Children who fail to improve in their nutritional status in the subsequent measurements (follow up) be referred to the next health institution for better management. For further details *refer to the core module unit 2, section 2.10*

### **2.11. Prevention & Intervention**

Give nutrition education to mothers or caregivers on **essential Nutrition Actions (ENA)** that need to be promoted are the following:

Exclusive breast-feeding up to 6 months of age

Complementary feeding at the age of 6 months with continued breast feeding up to 2 years and beyond

Continued feeding of the sick child( Importance of continued feeding during diarrheal attack or other illness)

Maternal nutrition: Increased consumption of balanced diet ~~during pregnancy~~ and lactation by mothers

Vitamin A supplementation for children 6-59 months and for lactating mothers within 8 weeks after delivery.

Iron and folic acid supplementation to





## INTRODUCTION

Appropriate feeding practices are important for the survival, growth, development, health and nutrition of infants and children everywhere. Every pregnant and lactating women needs adequate nourishment as means of attaining and maintaining health and recognizing that infant malnutrition is part of a wider problem .The health of infants and young children cannot be isolated from the health and nutrition of women. Therefore, Advising, and counseling women's are an important elements in the health care intervention. The following are important points with regard to micronutrient deficiency disorders.

### HOME TAKE MESSAGE WITH REGARD TO PRACTICE TO IMPROVE INFANT NUTRITION DURING THE FIRST SIX MONTHS.

#### 1. Initiate breastfeeding within one hour of birth.

Early initiation will be important for :

Take advantage of the newborns intense suckling reflex and alert state.

Stimulating breast milk production.

Serving as the baby's first immunizations. The infant will immediately benefit from the antibodies present in the colostrums (The first milk)

Minimizing maternal postpartum hemorrhage.

Fostering mother-child bonding.

#### 2. Establish good breast feeding skills (proper positioning, attachment, and effective feeding)

#### 3. Breast-feed exclusively for the first six months.

Breast milk should be the baby's first taste. There should be no prelacteal feeds such as water, other liquids, or ritual foods.

Breast milk completely satisfies an infant's nutritional and fluids needs for the first six-months.

Exclusively breastfed infants are at much lower risk of dying than the other children whom are on different feeding scheme.





## UNIT FOUR

### TASK AND ROLE ANALYSIS

Table 4.1. Knowledge objective and essential tasks of the health center team (health officer, Public health nurse, Medical laboratory technician and sanitarians)

	Learning objective (expected outcome)	HO	PHN	EH	MLT	Activity
Knowledge	Define and describe types of micronutrient deficiency	Define and describe types of micronutrient deficiency	Define and describe types of micronutrient deficiency.	Define and describe types of micronutrient deficiency.	Define and describe types of micronutrient deficiency.	Define and describe types of micronutrient deficiencies.
	List causes and risk factor of Micronutrient deficiencies.	List causes and risk factor of Micronutrient deficiencies.	List causes and risk factor of Micronutrient deficiencies.	List causes and risk factor of Micronutrient deficiencies.	List causes and risk factor of Micronutrient deficiencies	List causes and risk factor of Micronutrient deficiencies



	Describe the Magnitude and contribution of micronutrient deficiencies to over all childhood and adult health problems locally and nationally	Describe the Magnitude and contribution of micronutrient deficiencies to over all childhood and adult health problems locally and nationally	Describe the Magnitude and contribution of micronutrient deficiencies to over all childhood and adult health problems locally and nationally	Describe the Magnitude and contribution of micronutrient deficiencies to over all childhood and adult health problems locally and nationally	Describe the Magnitude and contribution of micronutrient deficiencies to over all childhood and adult health problems locally and nationally	Describe the Magnitude and contribution of micronutrient deficiencies to over all childhood and adult health problems locally and nationally
--	--	--	--	--	--	--

Table 4.2. Knowledge objective and essential tasks of the health center team (health officer, Public health nurse, Medical laboratory technician and sanitarians)

	Learning Objective (Expected Outcome)	HO	PHN	EH	MLT	Activities
Knowl edge	Describe the assessment Of micronutrient deficiency.	Describe the assessment Of micronutrient deficiency.	Describe the assessment Of micronutrient deficiency	Describe the assessment Of micronutrien t deficiency	Describe the assessment Of micronutrient deficiency	. Describe the assessment Of micronutrient deficiency
	Describe the principle and Treatment of micronutrient Deficiency.	Describe the principle and Treatment of micronutrient Deficiency.	. Describe the principle and Treatment of micronutrient Deficiency.	-----	----	. Describe the principle and Treatment of micronutrient Deficiency.
	Describe the pathogenesis Of micronutrient deficiency	Describe the pathogenesis Of micronutrient deficiency	---	---	---	Describe the pathogenesis Of micronutrient deficiency





<p>Believe in promoting proper Feeding of infants and balanced diet for adults.</p> <p>Believe in utilization of health services</p> <p>Both by children and adults in order to reduce the burden of micronutrient deficiency.</p>	<p>Advocate continued Feeding of a child and balanced diet for adults.</p> <p>Advice mothers, care takers</p> <p>And CHW to promote the utilization of existing health services in order to minimize micronutrient deficiencies.</p>	<p>Advocate continued Feeding of a child and balanced diet for adults.</p> <p>Advice mothers, care takers</p> <p>And CHW to promote the utilization of existing health services in order to minimize micronutrient deficiencies.</p>	<p>Advocate continued Feeding of a child and balanced diet for adults.</p> <p>Advice mothers, care takers</p> <p>And CHW to promote the utilization of existing health services in order to minimize micronutrient deficiencies.</p>	<p>Advocate continued Feeding of a child and balanced diet for adults.</p> <p>Advice mothers, care takers</p> <p>And CHW to promote the utilization of existing health services in order to minimize micronutrient deficiencies.</p>	<p>Advocate continued Feeding of a child and balanced diet for adults.</p> <p>Advice mothers, care takers</p> <p>And CHW to promote the utilization of existing health services in order to minimize micronutrient deficiencies.</p>	<p>-Educate mothers care give and (health extension workers, about the importance of balanced diet for children and adults to reduce micronutrient deficiencies.</p> <p>Teach about the importance of health services for children and adults to utilize the existing health services at the maximum possibility.</p>
--	--	--	--	--	--	---

	<p>Up hold the idea that micronutrient deficiency is caused by deficiency of nutrients.</p>	<p>Educate mothers, care givers and CHW how to prevent Micronutrient deficiencies.</p>	<p>Educate mothers, care givers and CHW how to prevent Micronutrient deficiencies</p>	<p>Educate mothers, care givers and CHW how to prevent Micronutrient deficiencies</p>	<p>Educate mothers, care givers and CHW how to prevent Micronutrient deficiencies</p>	<p>Educate mothers, care givers and CHW how to prevent Micronutrient deficiencies</p>
--	---	--	---	---	---	---

**Table. 4.4.** Practice objective and essential tasks of the health center team (health officer, Public health nurse, Medical laboratory technician and

	<p>Demonstrate how to do laboratory tests</p> <p>On micronutrient deficiencies.</p> <p>Demonstrate the preparation of dietary formula for the treatment micronutrient deficiencies to mothers and care givers.</p>	<p>Carry out laboratory investigation</p> <p>On micronutrient deficiencies</p> <p>Demonstrate the preparation of dietary formula for the treatment micronutrient deficiencies to mothers and care givers.</p>	<p>-----</p> <p>Demonstrate the preparation of dietary formula for the treatment micronutrient deficiencies to mothers and care givers.</p>	<p>-----</p> <p>Demonstrate the importance of clean water and utensils in the preparation of food in feeding with rich micronutrients</p>	<p>Carry out laboratory i</p> <p>Investigation pertaining to micronutrient deficiency.</p> <p>---</p>	<p>- Make a laboratory investigation on micronutrient deficiencies</p> <p>Show materials and ingredients to be used in the preparation and utilization of feeding formula in the treatment of micronutrient deficiency.</p>
--	--	---	---	---	---	---



Learning Objective (Expected out come)	HO	PHN	EH	MLT	Activities
Identify a case of micronutrients deficiency and demonstrate appropriate management.	Demonstrate the management principle, identify the complication and manage accordingly	Demonstrate appropriate feeding and drug administration and also  Provide proper nursing care to the clients.	-----	-----	Identify the case and its complication Mange the case by selecting appropriate treatment plan  Refer PRN -33(and )JTJ 62 0.72 ref



	<p>List causes and risk factors for pro Micronutrient deficiency.</p>	<p>List the different causes of micronutrient deficiency and their association with risk factors.</p>	<p>- Explain the cause of micronutrient deficiency in general and what risk behaviors are associated to it.</p>	<p>- Explain the relationship between the risk factors and development of micronutrient deficiency (HEW )          - Describe that micronutrient deficiency is caused by improper feeding, infection, diarrhea etc.</p>
--	---	---	---	---

Table 4.6. Attitude objectives and essential tasks of health extension workers and car givers

	Learning Objective (Expected out come)	CHW	Care giver	Activity
Attitude	Promote utilization of health service facilities for the treatment of Micronutrient deficiency.	Advise care givers to bring a child with Micronutrient deficiency the health service units to consult health worker	Advise friends and families to visit health worker the health service units in case of Micronutrient deficiency.	Educate care givers the importance of taking children with micronutrient deficiency to health d service institution  Encourage visits health service unit the case of Micronutrient deficiency.
	Advocate the importance of exclusive breast feeding in the first 4-6 months and continued feeding then after in reducing mortality and morbidity due to micronutrient deficiency.	Instruct mothers or care givers the importance of breast feeding in reducing morbidity and mortality from micronutrient deficiency..	Advise family friends and neighbors to continue breast feeding in a child with micronutrient deficiency.	Advocate / Promote breast feeding practices in prevention of micronutrient deficiency (HEW)  Encourage breast feeding practices of , family, in the prevention micronutrient deficiency.

Promote continued feeding of children with micronutrient deficiency. Advocate and encourage proper feeding of children with micronutrient deficiency by




Table 4.7. Practice objective and essential tasks of health extension workers and car givers.

	Learning Objective (Expected outcome)	HEW	Care giver	Activities
Practice	<p>Demonstrate preparation of feeding formula for the treatment of micronutrient deficiency and its proper use.</p> <p>Identify a case of micronutrient deficiency and demonstrate its appropriate management</p>	<p>Demonstrate preparation of feeding formulas their administration to the case of Micronutrient deficiency.</p> <p>Identify complications of Micronutrient deficiency and its degrees and advise the caregiver to feed the patient Properly.</p>	<p>Demonstrate properly how and what to prepare and administer to a child with micronutrient deficiency.</p> <p>Identify signs of symptoms of Micronutrient deficiency and its complications and decide whether there is a need for admission or referral</p>	<p>Show materials and ingredients to be used in the preparation of feeding formulas.</p> <p>Identify signs and symptoms of Micronutrient deficiency and administer proper feeding Practices (see the core module).</p>
	<p>Demonstrate proper communication to mothers or care givers pertaining to micronutrient deficiency.</p>	<p>Display Effective communication skill with mothers or care givers on treatment and prevention of Micronutrient deficiency.</p>		<p>Identify ways of educating Mothers/ care gives about micronutrient deficiency.</p>

## UNIT FIVE

### Glossary & Abbreviations

**Anthropometric assessment:** - Measurement of different body dimensions and proportions at different nutritional states and interpretation of the result by comparing to the standard to determine whether a person is malnourished or well nourished.

**Antioxidant:** - Micronutrients like vitamins A, C, E, & minerals, selenium which detoxify (scavenge) free radical species formed in the body and protect body cells from oxidative damage.

**Catch up growth:** - Rapid increase in weight and height of children after a period of nutritional deprivation in response to corrective dietary intervention.

**Communication:** the means of sending or receiving information, such as interpersonal communication, group discussion and mass communication.

**Folate:** a vitamin of the B complex found especially in leafy green vegetables, liver, and kidney.

**Fortification:** Addition of a nutrient in to a food that does not naturally contain it to increase the nutritive value of (food) by adding vitamins.

**Hygiene:** conditions or practices conducive to maintaining health and preventing disease, especially cleanliness.

**Latrine:** a toilet, especially a communal one used by individuals or communities.

**Malnourishment:** suffering from malnutrition, which may be under nutrition or over nutrition.

**Supplementation:** Periodic administration of a nutrient in the form of tablets, capsules or injections to an individual or a group of people to prevent the occurrence of deficiencies of the nutrient.

**Day Care Nutrition Rehabilitation Centers (DCNRC):** Feeding and nutrition demonstration centers attached to health units where mothers/care givers bring their

malnourished children and get them fed and see how to prepare balanced diet from locally available foodstuffs.

**Dehydration:** - Excessive loss of fluid and electrolyte from the body that impairs cellular function if not corrected timely.

**Emotional deprivation:** - State of mood change in a child that occurs following neglect of child (poor care given to the child by the mother or care giver).

**Exclusive breast-feeding:** - Breast-feeding of infants with no additional (supplementary) food or fluid administration.

**Recovery Syndrome:** - Fluid over load, congestive heart failure and death due administration of high protein and high caloric to a malnourished child during the acute (stabilization) phase of the management



**Weaning:** - Administration of food (solid or liquid including formula or cows milk) in addition to breast milk or without breast milk.



## UNIT SIX

### BIBLIOGRAPHY

#### **IODINE DEFICIENCY DISORDERS (IDD)**

- MOH-WNICEF. *The miracle of iodated salt: Ethiopia's commitment of universal salt iodations (USI), circular. 1995 No.9.*
- Luelseged T, Tefera B, Tesfaye B. *Iodine concentration of salt at household and retail shop level and knowledge attitude and practice of shop keepers and household food caterers, in Shebe Town sout*

- Kavisae P. Can Africa meet the goal of eliminating iodine deficiency disorders by the year 2000. *Bull food and Nutrition* 1996; 17(3): 262.
- Karmarkar MG, Pandav CS, Krishnamachari KAVR. Principle and procedure for iodine estimation: A laboratory manual. New Delhi, Indian Council of Medical Research, 1986.
- MOH. Micro Nutrient deficiencies control program in Ethiopia - profile draft (unpublished document 2001)
- Aboye and Urga K. Laboratory Manual. Ethiopian Nutrition Institute, Addis Ababa Ethiopia. (Unpublished document 1989).
- Demaeyer EM, Lowe Stein FW, Thilly CH. The control of Endemic Goiter. Geneva: WHO 1997.
- Anandk, Wajih SA, Prakashs and et al, A role for non-governmental organizations in monitoring the iodine content of salt in northern India. *Bull. WHO* 1995, 75(5): 71-75.

#### **IRON DEFICIENCY ANEMIA (IDA)**

- World Health Organization. Control of iron deficiency anemia in south-east Asia; (Report of an inter-country workshop) . Regional office for south-east Asia. New Delhi: World Health Organization; 1996.
- Tsinuel G, Tesfaye G, Tefera B, Sileshi T. The utility of pallor in detecting anemia in under five years old children. *Ethiopian Med. Journal*, 200; 38(2): 77-84.
- United Nation. Controlling iron deficiency. Geneva: world Health Organization; 1991.
- (ACC/SCN state-of-the-art series Nutrition policy Discussion Paper No.9 )  
Centre for disease control ((CDC). *Morbidity and mortality weekly report*. 1989; 38 (22): 400 - 4.
- Demaeyer E., Adeils -Tegman M. *World health statistics quarterly*. 1985; 38 (3) : 302-16.
- Demaeyer E.M., Dallman P., Gurney J.M., Sood SK., Srinikantia SG. Prevention and Control of iron deficiency anaemia through primary health care. Geneva: World health Organization [ WHO]; 1989.

- Lozoff B., Brittenham, G.M., Wolf A.W. et al. Iron deficiency anaemia and iron therapy. Effects on infant developmental test performance. *Paediatrics*. 1987; 79: 981-995
- Walter T. Infancy: Mental and motor development . *Am.J.Clin .Nutr*. 1989; 50: 655-66.
- Lozoff B., Brittenham G.M., Viteri F.E. et al. The effects of short -term iron therapy on developmental deficits in iron deficient anaemic infants. *J. Paediatrics*. 1982: 100:351-57. Walter T., De Andraca I., Chadud P., Perales C.G. Iron deficiency anaemia: Adverse effects on infant psychomotor development. *Paediatrics*. 1989; 84:7-17
- Pollitt E., Saco-Pollitt C., Leibel R.,I., Viteri F.E. Iron deficiency and behavioural development in infants and pre-school children. *Am.J. Clin. Nutr*. 1986;43:555-56
- Pollitt E., Leibel R.L., Greenfield D. Iron deficiency and cognitive test performance in pre-school children. *Nutr. Behav*. 1983; 1:137-46
- Soewondo S., Husaini M., Pollitt E. Effect

- Hussin M.A., Hassan H.A., Abdel-Ghaffar A.A., Salem S. Effect of iron supplementation on occurrence of diarrhoea among children in rural Egypt. *Food. Nutr. Bull.* 1988; 10(2): 35-39.



- J.L Diseases control priorities in developing countries. OUP, New York, 1993:421-51*
- *Boutry M., Needlman R. Use of diet history in screening iron deficiency anaemia. Paediatrics. 1996;96 (6Pt.1): 1138-42.*
  - *Halberg G.1 Bioavailability of dietary iron in man. Annu. Rev.Nutr. 1981:123-47*
  - *Lynch SR. Interaction of iron with other nutrients. Nutr. Rev. 1997;55 (4): 102-10*
  - *Lynch SR.Cook JD. Interaction of Vitamin C and iron. Ann. NY. Acad SCI 1980; 355:32-43.*
  - *Disler D.B., Lync, S.R., Charlton R.W. et al. Studies on the fortification of cane sugar with iron and ascorbic acid. Br. J.Nutr. 1980; 34:116-33.*
  - *Cook C.J., Monsen E.R. Vitamin C, the common colds and iron absorption. Am.J.Clin. Nutr, 1977;30:235-41.*
  - *Siegenberg D., Baynes R.D., Bothwell T.H., et al. Ascorbic acid prevents the dose dependent inhibitory effects of polyphenols and phytates on nonhaeme iron absorption Am.JClin.Nutr. 1991;53:357-41*
  - *Gillooly M., Bothwell T.H., Bothwell T.H., Torrance J.D. et al. The effect of organic acids, phytates and polyphenols on the absorption of iron from vegetables. Br.J.Nutr 1983;43: 271-9*
  - *Ballot, D.Baynes,R.D., Bothwell, T.H., Torrance J.D., et al. The effects of fruit juices and fruits on the absorption of iron from rieve meal. Brit.J.Nutr,57;1987:331-43*
  - *Derman, D.P., Bothwell, T.J., Torrance, J.D. et al. Iron absorption from maize (zeamays ) and sorghum (sorghum vulgare) beer. Brit, J.Nutr. 1980; 43:271-9*
  - *Cook J.D., Monsen E.R. Food iron absorption in human subjects. III, Comparison of effect of animal protein on nonhaeme iron absorption. Am.J.Clin . Nutr. 1976;29:859-67*
  - *Lynch S.R.,Hurrel R.F., Dassencto S.A., Cook, C.J. The effect of dietary proteins on iron bioavailability in man. Adv.Exp.Med.Biol. 1989; 249:117-32*
  - *Peter RG.et.al. Iron Bioavailability from diets by different socio-economic strata of the Venezuelan population.J.Nutr. 1995;125(7): 1860-8*



- Armstrong J.C., Chane T. Identification of hookworm species in Ethiopia. *Ethiop Med.J.* 1983; 21; 155-9
- Desalegn S. Prevalence of anaemia in pregnancy in Jimma town South-western Ethiopia. *Ethiop. Med.J* 1993; 31(4): 251-8.
- Ahmed zein Z. Haematocrit levels and anaemia in Ethiopian children. *East Afr. Med.J.* 1991; 68(6): 412-9.
- Stoltzfus R.J., Chwaya H.M., Albonico M., Schulze K.J., Salvioli L., Tielsch J.M. Serum ferritin, erythrocyte protoporphyrin and haemoglobin are valid indicators of iron status of school children in a malaria holoendemic population. *J.Nutr.* 1997; 127(2): 293-8
- Oski F.A. Iron deficiency in infancy and childhood. *N.Eng.Med.J.* 1993; 329(3): 190-3.
- World Health Organization. Prevention and management of severe anemia in pregnancy (report of technical working group). Geneva: World Health Organization; 1991. (WHO/FHE/MSM/93.5)
- World Health Organization. Indicators and strategies for iron deficiency and anaemia programmes (Draft report of the WHO/UNICEF/UNU consultation). Geneva: World Health Organization; 1994.

#### **VITAMIN A DEFICIENCY (VAD)**

- *Natl Acad Sci* 81, 7627-7631, -1.84-1 T.5 -1.73 Td(-)Tj/



carotene and lung cancer promotion in heavy smokers- a plausible relationship. *J Nat. Cancer Inst.* Nov. 6, 1996. 88(21), 1513-1515.

- The Alpha-tocopherol, Beta-carotene Cancer Prevention Study Group, The effect of vitamin E and beta-carotene on incidence of lung cancer and other cancers in male smokers. *N Engl J Med* 330, 1029-1035, 1994.

- Omenn GS, et al., The beta-carotene and retinol efficacy trial (CARET) for chemoprevention of lung cancer in high risk populations: *Smokersakevent35, nd5 19962al.*,

- oHkerBookRET)Nursiof roup,.0004 Tc59.0278 Tw 1.163

- Waterston T, editors. *Diseases of children in subtropics and tropics*. 4th ed. Edward Arnold, 1991.376-379.
- Aust-Kettis A, Bjomesjo E, Mannheimer E, Cvibah T, Clark P, Debele M. *Rickets in Ethiopia*. *Ethiop Med J* 1965;3:109-121.
- Eslietu M. *Analysis of pediatric admissions to Jimma hospital pediatrics ward: a 3 year retrospective study*. *Bull JIHS* 1994;4: 1-11.
- *Etiology of nutritional rickets: geographic variations (commentary)*. *J Pediatr* 1996; 128(5): 600-601.
- Kloos g Ahmed Z. *The ecology of health and disease in Ethiopia*. Westview press. Boulder, Sanfransisco, Oxford. 1993:96.
- 1 1. Wills XM Pbillips JB, Day RC, Bateman EC. *Phytic acid and nutritional rickets in inunigrants*. *Lancet* 1972; 1:771-773.
- 1"-). Kooh SW, Fraser D, Reilly BJ, et al. *Rickets due to calcium deficiency*. *N Engi J Med* 1977; 297:1264-66.
- Clenients M Johnson L, Fraser DR. *A new mechanism for induced vitamin D*

- Abe J, Nakano T, Nishii Y, et al., A novel vitamin D-3 analog, 22-2oxa-1,25-dihydroxyvitamin D-3, inhibits the growth of human breast cancer in vitro and in vivo without causing hypercalcemia. *Endocrinology* 1991; 129:832-837;
- Thomas M, Tebbutt S, and Williamson R: Vitamin D and its metabolites inhibit cell proliferation in human rectal mucosa and a colon cancer cell line. *Gut* 1992; 33:1660-1663.
- Latham, M.C., *Human nutrition in the Developing World*, Rome: F.A.O; 1997.
- Latham, M.C., *Human nutrition in tropical Africa*. Second edition, F.A.O, Rome; 1979
- Mahan LK., Stump SE. *Kruse's food, Nutrition and diet therapy*, 9<sup>th</sup> ed., USA: Saunders, 1996.

## ZINC DEFICIENCY

- Odeh M: The role of zinc in acquired immunodeficiency syndrome. *J Int Med* 1992; 231:463-469.
- McClain C, Stuart M, Vivian B, et al: Zinc status before and after zinc supplementation of eating disorder patients. *J Am Col N* 1992;11:694-700; and Varela P, Marcos A, Navarro M: Zinc status in anorexia nervosa. *Ann Nutr M* 1992;36:197-202.
- Tikkiwal M, et al., Effect of zinc administration on seminal zinc and fertility of oligospermic males. *Ind J Physiol Pharmacol* 31, 30-34, 1987; and Netter A, et al., Effect of zinc administration on plasma testosterone, dihydrotestosterone and sperm count. *Arch Androl* 7, 69-73, 1981.
- Goldenber RL, et al., The effect of zinc supplementation on pregnancy outcome. *JAMA* 274, 463-468, 1995.
- Eby GA, Davis DR, and Halcomb WW, Reduction in duration of common colds by zinc gluconate lozenges in a double-blind study. *Antimicrob Agents Chemother* 25, 20-24, 1984.



## **UNIT SEVEN**

### **ANNEXES**

#### **7.1 ANSWER KEYS**

##### **7.1.1 KEYS FOR THE CORE MODULE (ALL CATEGORIES)**

**Q.No.1. C**

**Q.No.2. E**

**Q.No.3. E**

**Q.No.4.A Marasmus**

**a. Kwashiorkor**

- b. Marasmic –kwashiorkor
- c. Underweight
- d. Stunting and wasting

Q.No.5. D

Q.No.6. Kwashiorkor

- B. Pitting edema
- C. Gray and easily pluckable hair
- D. Miserable and apathetic
- E. Loss of muscle & preservation of subcutaneous fat

**Marasmus**

- A. Loss of both subcutaneous fat and muscle (skin and bone appearance)
- B. Irritability and moodiness
- C. Wizen monkey faces (old man appearance)
- D. Absence of edema

Q.No.7.

- a. Anthropometric assessment
- b. Biochemical or laboratory, assessment
- c. Epidemiological (dietary assessment)

Q.No.8.

- d. Acute stabilization phase
- e. Rehabilitation phase

Q.No.9. a

condition called recovery syndrome)

Q.No.10. D

Q.No.11. D

Q.No.12. D

- Q.No.13. E  
Q.No.14. A  
Q.No.15. D  
Q.No.16. D  
Q.No.17. D  
Q.No.18. D  
Q.No.19. D  
Q.No.20. C  
Q.No.21. B  
Q.No.22. D  
Q.No.23. D  
Q.No.24. D  
Q.No.25. E

## 7.1.2.KEYS FOR SATELLITE MODULES (SPECIFIC PROFESSIONAL CATEGORIES)

### 7.1.2.1. HEALTH OFFICERS

- Q.No. 1. E  
Q.No. 2. D  
Q.No. 3. E  
Q.No. 4. A. Goiter with its complications  
B. Mental retardation and poor academic performance  
C. Poor physical work out put  
D. Poor Physical growth  
Q.No. 5. D  
Q.No. 6. E  
Q.No. 7. D

- Q.No. 8 E
- Q.No. 9. D
- Q.No. 10. D
- Q.No. 11. E
- Q.No. 12. E
- Q.No. 13. D
- Q.No. 14. E
- Q.No. 15. E
- Q.No. 16. B
- Q.No. 17. E
- Q.No. 18. E
- Q.No. 19. E
- Q.No. 20. B
- Q.No. 21. E
- Q.No. 22. E
- Q.No. 23. A
- Q.No. 24. A
- Q.No. 25. E
- Q.No. 26. E
- Q.No. 27. E
- Q.No. 28. E

### 7.1.2.2. PUBLIC HEALTH NURSE

Key for Public Health Nurse

1. Antenatal care  
Delivery  
Post-natal care/FP  
Immunizations  
Well baby clinic  
Sick baby clinic

2. D.

3.

Food diversification

Food fortification

Vit A supplementation

4. A

5. D.

6. C.

7. D.

8. 150 microgram per person

9. D.

10. Beef, cheeks, fish, Lamb, liver, and pork.

11.

Prolonged breast feeding with out complementary feeding.

Heammorhage.

Infections

Low consumption (Dietary)

Repeated pregnancy.

13. "Dry Eye" in Greek.

Function as anti oxidant

Maintain normal taste and smell

Promotes normal growth and development

Aids wound healing

Promotes normal fetal growth

Helps synthesize DNA and RNA.

Promotes cell division cell repair cell growth

Maintain normal level vit A in blood.





- B. Personal, environmental and food hygiene
- C. Importance of taking their children to the health institutions for Growth monitoring
- D. Importance of getting their children immunized
- E. Stimulation and proper treatment of children

**7.2. Recommended Dietary Allowance (RDA) For selected micronutrients:  
Estimate of adequate daily intake by the Food and Nutrition Board of the National  
Research Council, 1989 by age group.**

**FOLIC ACID**

**Age RDA**

- 0-6 months 25mcg
- 6-12 months 35mcg
- 1-3 years 50mcg
- 4-6 years 75mcg
- 7-10 years 100mcg

**Males**

- 11-14 years 150mcg
- 15+ years 200mcg

**Females**

- 11-14 years 150mcg
- 15+ years 180mcg
- Pregnant 400mcg
- Lactating 1st 6 mos 280mcg
- 2nd 6 mos 260mcg

## VITAMIN A

### **Age Retinol Equivalents International Units**

0-6 months 375RE 2,100IU

6-12 months 375RE 2,100IU

1-3 years 400RE 2,000IU

4-6 years 500RE 2,500IU

7-10 years 700RE 3,300IU

### **MALES**

11+ years 1,000RE 5,000IU

### **FEMALES**

11+ years 800RE 4,000IU

Pregnant 800RE 4,000IU

Lactating 1st 6 mos 500RE +2,500IU

2nd 6 mos 500RE +2,500IU

## VITAMIN D

### **Age RDA**

0-6 months 7.5mcg

6-12 months 10mcg

1-10 years 10mcg

### **MALES**

11-18 years 10mcg

19-24 years 10mcg

25+ years 5mcg

### **FEMALES**

11-18 years 10mcg

19-24 years 10mcg

25+ years 5mcg

Pregnant 10mcg

Lactating 10mcg

## IODINE

Age	RDA
0-6 months	40mcg
6-12 months	50mcg
1-3 years	70mcg
4-6 years	90mcg
7-10 years	120mcg
11+ years	150mcg
Pregnant	375mcg
Lactating	200mcg

## IRON

Age	RDA
0-6 months	5mg
6-12 months	10mg
1-3 years	10mg
4-6 years	10mg
7-10 years	10mg

---

**MALES**

11-18 years	12mg
19+ years	10mg

**FEMALES**

11-50 years	15mg
51+ years	10mg
<b>Pregnant</b>	30mg
<b>Lactating</b>	15mg

---

**Zinc**

---

Age:	RDA:
------	------

---

0-6 months	5mg
6-12 months	5mg
1-10 years	10mg

**Males:**

11+ years	15mg
-----------	------

**Females:**

11+ years	12mg
-----------	------

---

**7.3. THE AUTHORS****TEFERA BELACHEW (MD, MSC, DLSHTM),**

Associate professor, Population and Family Health Department, Jimma University. He obtained his degree of doctor of medicine from the former Jimma Institute of Health Sciences and Master of Science in human nutrition from University of London, London School of Hygiene and Tropical Medicine, UK.

**ABEBE GEBREMARIAM (BSC, MPH)**

Associate professor, Population and Family Health Department, Jimma University. He obtained his BSc in public health from the former Gondar college of Public Health and his MPH from Tulane University, School of Public health and Tropical Medicine, USA.

**WORKU LEGESSE(BSC, MSC, PhD),**

Assistant professor in school of Environmental Health, Jimma University. He obtained his Bachelor degree in biology from Asmara University and his MSc in environmental Sanitation from Ghent University Belgium and His PhD in ecology from Cork University in the Republic of Ireland.

**TSEGAYE ASRES (BSC, MSC, DLSHTM),**

Lecturer in the School of Nursing, Jimma University. He obtained his BSc in nursing from the former Jimma Institute of Health Sciences and His MSc from University of London, London school of Hygiene and tropical Medicine, UK.

**TAMRAT TIGABU (BSC, MSC),**

Lecturer in the School of Nursing, Jimma University. He obtained his BSc in nursing from the Gujarat University India his MSc in nursing from The Queen's University of Belfast, Northern Ireland.

**SEID TIKU (BSC),**

Graduate assistant II in the school of Environmental Health, Jimma University. He obtained his BSc in environmental health from Jimma University.

**ESTIFANOS KEDEDE (BSC, M.MED.SC)**

Lecturer in the School of Medical Laboratory Technology, Jimma University. He obtained his BSc in Medical Laboratory Technology, form Jimma University and his M.Med.Sc from University of Free State, Republic of South Africa.

**GETACHEW DEMEM (MD)**