

7 KLV DUWLFOH LV UHSULQWHG RQ WKH &DUWHU &HQWHU¶V ZHEVLWH  
*Tropical Medicine and Hygiene.*

Am. J. Trop. Med. Hyg., 83(3), 2010, pp. 534-541

doi:10.4269/ajtmh.2010.10-0033;

Copyright © 2010 by [The American Society of Tropical Medicine and Hygiene](#)

## **Effects of Annual Mass Treatment with Ivermectin for Onchocerciasis on the Prevalence of Intestinal Helminths**

**Julie Gutman\***

# INTRODUCTION

twice per year IV therapy in northeastern Ecuador on STH. However, no studies have been done to look at the effect on the prevalence of STH following annual mass drug administration (MDA) with IV for onchocerciasis that occurs at approximately 40 million treatments per year in Africa. <sup>24</sup>

The use of annual IV distribution to prevent morbidity caused by onchocerciasis began in Imo state, Nigeria, in 1993 and reached statewide in 1995 following a national onchocerciasis assessment survey that showed high prevalence of onchocerciasis throughout much of southeastern Nigeria. <sup>6, 25</sup> This program was begun as a combined effort of the State Ministry of Health, the Lions Clubs, and the River Blindness Foundation. For a time it also received support from the African Program for Onchocerciasis Control (APOC). The distribution program is now run by the State Ministry of Health in conjunction with The Carter Center. Imo State is composed of 27 districts known as local government areas (LGAs). On the basis of the 1995 disease mapping for onchocerciasis, 18 of the 27 LGAs in Imo state receive annual IV therapy because onchocerciasis is a public health problem there. In affected villages,

**HO**

## METHODS

**Sampling methodology.** From July to August, 2008, we performed a cross-sectional survey of STH prevalence in Imo state, Nigeria. We used a stratified sampling procedure to choose 40 villages, 20 treated and 20 untreated villages (

his/her name recorded. The child or child's guardian was instructed to put a walnut sized amount of feces (size showed with a rock) of his or her stool from the next morning into the collection cup using a wooden stick, which was provided. The child/guardian was also instructed to wash his/her hands

skewed toward low egg counts. For this reason, logarithmic transformation and geometric means were calculated for fecal egg density using  $\text{antilog}-1$  where  $x = \text{number of}$



untreated area ( $P = 0.07$ ). Only light intensity *Trichuris* infections were seen in the treated areas, whereas 3 children (5.7%) had moderate intensity infections in the untreated area ( $P = 0.55$ ). Heavy infections with hookworm occurred in 3.2% and 1.8% of children in the treated and untreated area, and moderate intensity infections occurred in 2.1% and 2.7% of children in the treated and untreated area ( $P = 0.77$ ). Only one child was found to be infected with *S. mansoni*, none were infected with *Schistosoma hematobium*.

**Preschool-aged.** Among preschool-aged children, who had never received IV even in treated areas, there was a significant difference in the prevalence of infection with *Trichuris* in the treated (1%) compared with untreated areas (8%) ( $P = 0.019$ ). The difference in the prevalence of *Ascaris* infection approached statistical significance (3% versus 10%,  $P = 0.051$ ). There was no difference in the prevalence of hookworm infection (21% versus 27%,  $P = 0.30$ ). Geometric mean egg counts per gram of stool among all children 2–4 years of age were higher in untreated versus treated MDA; this difference was significant for *Ascaris* and *Trichuris* but not for hookworm (*Ascaris*: 0.91 [95% CI 0.30–2.83] versus 0.24 [95% CI 0–0.59],  $P = 0.04$ ; *Trichuris*: 0.39 [95% CI 0.12–1.14] versus 0.05 [95% CI 0–0.14],  $P = 0.01$ ; hookworm: 2.44 [95% CI 1.30–4.14] versus 1.74 [95% CI 0.82–3.12],  $P = 0.36$ ). The range of eggs per gram of stool in untreated versus treated areas was 0–8,648 for *Ascaris*, compared with 0–6,000 in the treated region, 0–12 versus 0–6 for *Trichuris*, and 0–24 versus 0–1,632 for hookworm. Three children had *Ascaris* infection of moderate intensity; one in the treated and two in the untreated areas, respectively ( $P = 1$ ). The infections with *Trichuris* and hookworm were all of light intensity. There was one case of infection with *S. mansoni*.


**Effects of ivermectin on the need for community treatment.** From a community perspective, nearly all the villages required at least school-based therapy for hookworm;





**View larger version (23K):**  
[\[in a new window\]](#)

FIGURE 2.



**View this table:**  
[\[in a new window\]](#)

TABLE 3  
Univariate and multivariate analysis of factors contributing to  
infection among SACs



untreated groups. Although we found a significant impact of ivermectin on the prevalence of *Ascaris* and *Trichuris*, there was no difference in the prevalence or intensity of infection with hookworm in the treated versus untreated areas. The lack of efficacy against hookworm, the STH most associated with causing anemia and intestinal inflammation, explains the lack of difference in growth parameters.

infection, we cannot be sure that the treated villages had the same initial prevalence and intensity of STH as the untreated villages. However, given the similar hookworm findings between areas, and the known properties of ivermectin against the STHs, we think our assumption that *Ascaris* and *Trichuris* levels were similar before IV interventions began is reasonable.

Further studies are indicated to determine if annual IV and albendazole will suffice to control STHs in IV-treated onchocerciasis areas where STH prevalence is > 50%, so averting the need to invest in an additional treatment round. In addition, preschool-aged children are at risk for STH infections and better studies and age appropriate drug preparations are needed to guide recommendations for STH treatment in this age group.

---

Received January 18, 2010. Accepted for publication May 5, 2010.

## ACKNOWLEDGMENTS

We thank Josephine Obiezu who entered the data; Stanley Asiawuchi, Afam Okonkwo, and Francis Ugwu who prepared and read all the slides; Jon Nnachi and Aaron Iroaja who assisted in data collection; Aryc Mosher for preparing the map in Figure 1; all the community health workers, villagers, and children who participated; and Daniel Newman for proofreading and allowing his wife (Julie Gutman) to leave him for a month of field work.

Financial support: We gratefully acknowledge the Izumi Foundation and the Bill and Melinda Gates Foundation for their generous funding. Julie Gutman's salary was supported in part by PHS Grant UL1 RR025008 and KL2 RR025009 from the Clinical and Translational Science Award program, national Institutes of Health, National Center for Research Resources and a fellowship grant from the Pediatric Infectious Disease Society [PIDS].

Authors' addresses: Julie Gutman, Emory University School of Medicine, Department of Pediatric Infectious Disease, E-mail: gutmanjr@gmail.com. Emmanuel Emukah, Njideka Okpala, and Andrew Obasi, The Carter Center Southeast Programs, Nigeria, E-mails: emukahe@yahoo.com, talk2nj2003@yahoo.com, and andyobasi2@yahoo.com Chinyere Okoro, Parasitology Unit, Microbiology Department, Federal Medical Center, Owerri, Imo State, Nigeria, E-mail: ihuarulam2002@yahoo.com

## REFERENCES

1. Kirwan P, Asaolu S, Molloy S, Abiona T, Jackson A, Holland C, 2009. Patterns of soil-transmitted helminth infection and impact of four-monthly albendazole treatments in preschool children from semi-urban communities in Nigeria: a double-blind placebo-controlled randomised trial. *BMC Infect Dis* 9: 20.[CrossRef][Medline]
2. Chan M, 1997. The global burden of intestinal nematode infections <sup>2</sup> fifty years on. *Parasitol Today* 13: 438 -443.[CrossRef][Web of Science][Medline]
3. Kirwan P, Asaolu SO, Abiona TC, Jackson AL, Smith HV, Holland CV, 2009. Soil-transmitted helminth infections in Nigerian children aged 0 -5 months. *J Helminthol* 83: 261 -266.[CrossRef][Web of Science][Medline]
4. Uneke C, Eze KO, Oyibo PG, Azu NC, Ali E, 2007. *Soil-Transmitted Helminth Infection In School Children In South-Eastern Nigeria: The Public Health Implication*. The Internet Journal of Third World Medicine. Volume 4.
5. World Health Organization, 2002. *Prevention and Control of Schistosomiasis and Soil Transmitted Helminths: report of a WHO expert committee*. World Health Organization Technical Report Series 912.
6. Hopkins D, Eigege S, Miri ES, Gontor I, Ogah G, Umaru J, Gwomkudu CD, Matha W, Jinadu MY, Amadiogwu S, Oyeneka OK, Korve K, Richards F, 2002. Lymphatic filariasis elimination and schistosomiasis control in combination with onchocerciasis control in Nigeria. *Am J Trop Med Hyg* 67: 266 -272.[Abstract]
7. Emukah E, Oshuoha E, Miri ES, Onyenama J, Amazigo U, Obijuru C, Osuji N, Ekeanyanwu J, Amadiogwu S, Korve K, Richards FO, 2004. A longitudinal study of impact of repeated mass ivermectin treatment on clinical manifestations of onchocerciasis in Imo State, Nigeria. *Am J Trop Med Hyg* 70: 556 -561.[Abstract/Free Full Text]
8. Bleakley H, 2003. Diseases and development: evidence from the American South. *J Eur Econ Assoc* 1: 376 -386.[CrossRef]
9. Stephenson L, Latham MC, Ottesen EA, 2000. Malnutrition and parasitic helminth infections. *Parasitology* 121: S23 -S28.[CrossRef][Web of Science][Medline]





severely affected population. *Bull World Health Organ* 82: 563-571.[Web of Science][Medline]

21. Tatischeff S, Kebede A, Bulto T, Werkeneh W, Tilahun D, 1994. Effect of ivermectin (Mectizan) on intestinal nematodes. *Ethiop Med J* 32: 7-15.[Web of Science][Medline]

22.

31. Oberhelman RA, Guerrero ES, Fernandez ML, Silio M, Mercado D, Comiskey N, Ihenacho G, Mera R, 1998. Correlations between intestinal parasitosis, physical growth, and psychomotor development among infants and children from rural Nicaragua. *Am J Trop Med Hyg* 58: 470 -475.[Abstract]
32. Moore SR, Lima AA, Conaway MR, Schorling JB, Soares AM, Guerrant RL, 2001. Early childhood diarrhoea and helminthiases associate with long-term linear growth faltering. *Int J Epidemiol* 30: 1457 -1464.[Abstract/Free Full Text]
- 33.

Hoa Binh province, Vietnam. *Trans R Soc Trop Med Hyg* 103: 237 ± 241.[CrossRef][Web of Science][Medline]

41. Asaolu S, Ofoezie IE, Odumuviwa PA, Sowemimo OA, Ogunniyi TA, 2002. Effect of water supply and sanitation on the prevalence and intensity of *Ascaris lumbricoides* among pre-school-age children in Ajebandele and Ifewara, Osun State, Nigeria. *Trans R Soc Trop Med Hyg* 96: 600 –604.[CrossRef][Web of Science][Medline]
42. Chongsuvivatwong V, Pas-Ong S, McNeil D, Geater A, Duerawee M, 1996. Predictors for the risk of hookworm infection: experience from endemic villages in southern Thailand. *Trans R Soc Trop Med Hyg* 90: 630 –633.[CrossRef][Web of Science][Medline]