Acknowledgements

The Carter Center's Trachoma Control Program would like to acknowledge the support of numerous donors who have made the 2014 activities reviewed in this document possible:

Abbott Laboratories

Anonymous (4)

Robert and Joan Blackman Family Foundation

Blum Family Foundation

The Community Foundation for Greater Atlanta, Inc.

Crabby Beach Foundation

Mr. and Mrs. Pete Docter

Acronyms

API	Active Pharmaceutical Ingredients (specific to Pfizer Inc)
BCC	Behavior Change Communication
CHW	Community Health Worker

Indices for Ultimate Intervention Goals (UIGs)*

Surgery $$\frac{\text{Sum of surgeries to date}}{\text{Sum of surgeries to date + most recent backlog}}$$

Angelia Sanders, Associate Director of The Carter Center's Trachoma Control Program, proposed several new indicators in her presentation, "Facial Cleanliness and Environmental Improvement Indicators: Showing Impact Through Purpose". Ms. Sanders challenged the current indicators, explaining that while they tell the trachoma community that health education is being done and latrines are being constructed, they do not provide information about the level of comprehension of messages and the degree of latrine usage.

SAFE in Ethiopia

Presented by Mr. Oumer Shafi, NTD Team Leader, Federal Ministry of Health, Ethiopia

Background

The National Survey on Blindness, Low Vision, and Trachoma conducted in 2006 revealed that 2.8 million people in Ethiopia had low vision and 1.2 million people were blind. It was estimated that 87 percent of blindness was from avoidable diseases. The survey revealed that active trachoma was endemic in virtually all regions of the country, with more than 1.3 million people in the country living with TT. The results of the survey showed that Ethiopia had approximately 30 percent of the burden of trachoma in Sub-Saharan Africa.

There is growing momentum on the issue of Neglected Tropical Diseases (NTDs) in Ethiopia. In 2013, a national NTD master plan was launched, with regional states preparing their own NTD master plans. A NTD team was formed within the Federal Ministry of Health (FMOH) and NTD indicators became part of the national health management information system. National treatment registers and health extension worker (HEW) pocket guidelines are in the process of being developed, with plans for NTDs to be integrated into the existing health system. In order to better understand the NTD burden in Ethiopia, mapping of diseases that can be treated with preventative chemotherapy (PCT), such as lymphatic filariasis, schistosomiasis, and soil-transmitted helminthiasis, was conducted. The results from the Global Trachoma Mapping Project (GTMP) will further assist with the NTD efforts related to trachoma elimination.

Timeline of Events

2001: National guideline for Primary Eye Care developed[1]
2006: National guideline for mass antibiotics distribution developed; national task force for trachoma control established
2006-2007: Amhara region's baseline survey at zonal level
2008: Trachoma Campaign, formerly MalTra, launched[2]
2012: National Trachoma Action Plan (TAP) was prepared
2010-2014: GTMP and trachoma impact surveys conducted in Amhara region
2013: Trachoma becomes part of national NTD program under disease prevention and control directorate
2020: Target date for elimination

[1] A five-year document, currently in its 3rd cycle.
[2] MalTra (Malaria and Trachom3(n)4(d)-g-2(c)-4(u)eear 131.t4( )](n0 g 0 G [( )] TJd2)eea [. /P n>> 1 .

Table 1. ProgramA

Environmental Improvement (E)

Improved latrines—a latrine which includes a handwashing facility, a slab, and ventilation pipe—are in 6.76 percent of households in the Oromiya region. A total of 42.4 percent of households in the Amhara region have a latrine according to the DHS conducted in 2012–2014. Since 2009, 79 percent of the household latrine construction UIG has been achieved. In 2014 with the construction of 1,732,049 latrines, Ethiopia reached 86 percent of its annual construction goal of two million latrines. According to the 2014 Ethiopian Mini Demographic Health Survey, 57 percent of rural households own improved toilet facilities, and only 55 percent of total households have access to an improved source of drinking water.

In 2014, The Ethiopian Ministry of Health (MOH) revised its TAP based on the completed GTMP. The revised TAP describes a plan for the inclusion of the F and E components of the SAFE strategy in the Health Extension Package of the National Health Care System. To identify coverage of F and E components, the FMOH, through the TAP, identified water, sanitation, and hygiene (WASH) partners' locations and assumed that F and E activities were active in these districts. However, the extent of WASH partners' F and E activities or the extent to which they apply to trachoma elimination remains unclear.

Also in 2014, the Ethiopian government created a national, multisectoral WASH collaboration between governmental sectors—Ministry of Education, FMOH, Ministry of Finance and Economic Development—and donors called One WaSH. In addition to advocating for a consolidated WASH account, the initiative outlines sanitation and hygiene promotion in rural, urban, and institutional divisions depending on specific criteria. The Ethiopian government is providing about 10 percent of the funding for this 4.84-billion-dollar initiative. Determining points of collaboration and alignment between trachoma elimination objectives and the One WaSH program will be a priority for the program in 2015.

Programmatic Challenges:

More than 250





Table 1. Program Achievements in 2018

Indicator	UIG	National		Carter Center Assisted	
		Target	Achieved	Target	Achieved

Antibiotic Therapy (A)

In 2014, among children

Recommendation 2: All programs should consider conducting a brief survey to assess MDA coverage after every drug distribution.

Since MDA was not conducted in 2014, this recommendation was not applicable.

Recommendation 3: Whenever possible, national trachoma programs should publish data documenting their experiences related to levels of TF, stopping of MDA, follow up, and sampling frames used to assess level of TF.

Not completed. Planned for 2015.

Recommendation 4: All country programs should consider developing detailed plans to increase TT surgical output and quality to present at next year's program review meeting.

Completed.

Recommendation 5: The program should resume activities in the North, security permitting.

Insecurity continues to prevent activities in the Kidal and Gao regions.

Recommendation 6: Conduct cross-border collaborative meetings and plans by both the implementing partners and member states.

Completed. Trachoma program officials from Niger and Mali participated in each other's program reviews. For a week in the summer, the Malian program and its partners visited the field in Niger to observe trachoma control interventions.

Recommendation 7: National Programs should consider strengthening their system of mobilization and

Mali: TF Prevalence among Children 1-9 years

Baseline, 1996



2014




Mali
N
W
E
S
Timbuctou
Kidal
Mauritania
Niger
Kayes
Ségou

Table 1. Program Achievements in 2014

Indicator	UIG	National		Carter Center-Assisted	
		Target	Achieved	Target	Achieved
# of persons operated	29,887	15,000	17,745 (118%)	11,000	14,790 (134%)
# of women operated			N/R [5]		8,858 (60%)
# of surgeons trained		40	44 (110%)	30	30 (100%)
# of surgeons retrained			0 (0%)		0 (0%)
# of surgeons certified		40	40 (100%)	20	20 (100%)
Doses of azithromycin distributed during MDA	4,768,480	4,768,480	N/R	N/A [6]	N/A
Doses of tetracycline distributed during MDA	170,000	170,000	N/R	170,000	N/R
# of villages with health education	634	634	647 (102%)	634	647 (102%)

For supervision, a set allowance of days is given for supervisors at each level to go into the field; often this allowance does not account for the realistic amount of time to carry out supervision activities. The supervisor inspects training, introduction of drugs and other inputs. The community distributors noted the number of doses distributed in a disease-specific check sheet, where gaps of information at times encountered due

The program will meet with village leaders and district heads. The program has experienced a relative standstill in the dissemination of facial cleanliness messages in schools; as a result, the program wants to target academic advisors who train teachers to train one teacher per school on facial cleanliness messages and to include training on facial cleanliness m γ i ⁎ ȳ inclm on d] w φ d . γ t l’ n ơ

Niger: TF Prevalence among Children 1-9 years

Baseline, 2006-2007



2014



Niger: Prevalence of TT among Adults ≥ 15 years

Baseline, 2006-2007



2014




Algeria
< 500
Dosso


Niger
< 80

Tahoua
Tillaberi
Dosso
Burkina Faso
Benin

SAFE in Nigeria

Presented by Dr. Falam J. Nimzing, Trachoma Officer, The Carter Center, Nigeria and
Mr. Nicholas Olobio, Data Manager, National Trachoma Control Program, Federal Ministry of Health, Nigeria

Background

The Nigeria Blindness and Low Vision Survey of 2005-2007 noted that the overall prevalence of blindness in Nigeria was 0.79 percent and that the prevalence varied across the different geopolitical zones of the country (range: 2.8-6.1 percent). The survey identified cataract as the main cause of avoidable blindness in Nigeria, followed by trachoma. Trachoma is among the top 10 priorities in the country and part of the NTDs division of the FMOH.

Though eye care activities have been conducted in Nigeria since 1991 under the National Program for Prevention of Blindness (NPPB) in 2001 the National Trachoma Control Program was established in order to better focus on trachoma control activities. Regular coordination meetings are held for government and partners in trachoma control. With the support of ENVISION, a project funded by the United States Agency for International Development (USAID) and implemented by RTI International, the meeting for the development of a TAP was held in Abuja from June 24-28, 2013. The Nigerian TAP sets out the implementation details of the SAFE strategy in Nigeria with the goal of eliminating blinding trachoma by 2018.

In 2013, the GTMP and ENVISION supported household surveys to generate updated data on active and blinding trachoma and access to sanitation and safe water. In 2014, GTMP completed mapping of all local government areas (LGAs) in the country suspected of being endemic for trachoma, excluding any insecure areas. The Carter Center assisted with the mapping in five states in the Southeast. Currently, only Adamawa and Borno states remain to be mapped with assistance from other partners

Table 1. Program Achievements in 2014 N/A

Indicator	UIG	National		Carter Center-Assisted	
		Target	Achieved	Target	Achieved
# of persons operated	294,821	40,700	7,905 (19%)	N/A [7]	N/A
# of women operated	[redacted]	19,188	3,680 (19%)	N/A	N/A
# of surgeons trained	[redacted]	12	7 (58%)	N/A	N/A
# of surgeons retrained	[redacted]	N/A	N/A	N/A	N/A

SAFE Strategy, drug distribution, reporting, and mobilization. In some areas, the National Program integrated trachoma MDA training with other NTD MDA trainings. To enhance community sensitization, the program used posters and handbills. Radio stations also aired radio jingles in local languages to provide the benefits and dates of MDA. The program continues to meet with community, opinion, and religious leaders and schoolteachers to increase mobilization.

MDA supervision is carried out in three divisions. Pre-MDA supervision occurs during training. During the MDA, intra-MDA supervision ensures proper health education, dosing, and direct observed treatments compliance. In addition to conducting spot checks and coverage surveys, post-MDA supervision monitors timely reporting and proper disposal of containers. Nigeria's MDA reporting flows from communities, to Frontline Health Facilities, to LGA's, to states, to zones, and then to partners and FMOH.

Facial Cleanliness (F)

In 2014, the program conducted health education in 7,755 villages, achieving 64 percent of its target. The Carter Center assisted in reaching 855 of these villages. The National Program produced a Trachoma Instructional Program to sustain the progress already made on the elimination targets in Nasarawa and Plateau states. To implement this program, Nigeria held teacher workshops for two teachers from each school in the states of Nasarawa and Plateau, where the intervention took place. The National Program produced health education materials to aid teachers. In addition to providing health education in places of worship, market places, ceremonies, and in clinics, the National Program also aired radio jingles and TV life presentations.

Environmental Improvement (E)

The program constructed a total of 130 latrines, achieving 29 percent of its target. Until 2009, The Carter Center assisted all National Program latrine construction. With the exception of one partner's work, no official program-supported latrine construction took place in Nigeria in 2014. The National School Health Curriculum for trachoma included education on sanitation and hygiene and used materials such as posters, flipcharts, and billboards to help disperse these messages. In the past year, the National Program provided plastic containers for water in each classroom and provided wells or boreholes at home and schools. Sanitation officers, town announcers, and local government integration health teams helped promote sanitation and hygiene. The National Program encouraged communities to dig wells, rehabilitate water points, and harvest rain water. The government at state, LGA, and community levels collaborates with WASH institutions to promote access to water.

Programmatic Challenges

Political interests and lack of reporting by partners contribute to weak linkages and poor collaboration for the F and E components of the SAFE strategy. Partners and the Nigerian government's low commitment to F and E activities pose a challenge to water and sanitation improvement. Security challenges prevent trachoma interventions in some p

Targets for 2015 and Plans to Meet Targets

Surgery (S)

- Operate on 9,465 patients
- Train and certify 235 surgeons
- Retrain 30 surgeons

In order to meet the surgical LF elimination goal by the 2018 elimination date, Nigeria will standardize the training curriculum to ensure that all TT surgeons have demonstrated an accepted level of competency. According to the work plan for clearing the TT surgery backlog by 2018, the target of operating 9,465 patients is in addition to the 31,980 surgeries set out by the plan to be performed in 2015.

Antibiotic Therapy (A)

- Distribute 8,675,865 doses of azithromycin
- Distribute 173,517 doses of tetracycline

In 2015, the program will continue its current activities for MDA mobilization. The National Program has placed FMOH staff in zonal offices who will aid in collating data from the states and transmit the data to the federal level. Additionally, the program will introduce an integrated NTD reporting form, developed through the support of RTI International.

Table 1. Program Achievements in 2014

Indicator	UIG	National		Carter Center-Assisted	
		Target	Achieved	Target	Achieved
# of persons operated	121,062 (28 of 79 districts)				

Recommendation 4

Western Equatoria

Chad
Central African Republic


Jonglei


Jonglei

SAFE in Sudan

Presented by Dr. Nabil Aziz Mikhail, Country Representative, The Carter Center, Sudan
Zeinab Abdalla, Program Officer, The Carter Center, Sudan
Dr. Balgesa Mohammed, National Coordinator, Trachoma Control Program, Federal Ministry of Health, Sudan

Background

The FMOH has been working towards trachoma control since 1962, when trachoma was incorporated into the NPPB. The Academy of Medical Sciences and Technology took over the leadership of the program in the 1990s as contractor on behalf of the FMOH. In 2005, the FMOH relocated the Trachoma Control Program to the NPPB. The elimination of blinding trachoma is one of the FMOH's priorities and government funds are allocated to support the program. In 2012, the government allocated 1.5 million USD for five years to help support The Carter Center's partnership for trachoma control. There is a strong coordination mechanism between the government

In 2014, in order to increase awareness about and during MDA, the program mobilized politicians and

Status of 2014

Sudan:TF Prevalence among Children 1-9 years

Baseline, 2006-2010



2014

Sudan: Prevalence of TT among Adults ≥ 15 years

Baseline, 2006



Presented by

Table 1. ProgramAchievements in 2014

Indicator	UIG	National

schools and communities. Radio announcements, sports events, and television talk shows were conducted and public announcements were made to promote MDA mobilization.

For MDA supervision, the DOH first makes courtesy calls on

Recommendation 2:

will also develop a six-month post-operative follow-up strategy and will organize TT surgery camps closer to settlements. Disaggregating the data by gender, the National Program will standardize data collection tools surgical logbooks, TT patient registers, TT case finders' registers, and summary reporting forms—and will pilot these tools, tracking their success through a new eHEALTH reporting system. To increase surgery



South Sudan
Rwanda

Table 1. Summary of National Data from Trachoma Control Programs (Carter Center Assisted Countries)
National Data as Reported for 2014 at the Sixteenth Annual Program Review, Atlanta, Georgia, March 2-4, 2015

	Mali	Niger	Sudan

Table 2. National Trachoma Control Program Annual Targets 2015 (Carter Center-Assisted Countries)

Targets[†] as Presented at the Sixteenth Annual Program Review, Atlanta, Georgia, March 2-4, 2015[§]

Table 3. Carter Center-Assisted Implementation of SAFE (Carter Center-assisted outputs)**

Summary of Interventions per Country, January - December 2014

Indicators	Mali	Niger	Sudan	South Sudan	Ethiopia-Amhara	Nigeria	Total
Surgery							
Persons operated for trichiasis	2,742	14,790	1,924	213	40,450	0	60,119
2014 Target	6,000	11,000	2,000	N/A*	61,537		




2,490
634
651
10,850
12,100
27,107

Where do opportunities for behavioral change communication exist?

Presented by Tara Brant, BCCE Consultant, The Carter Center, Ethiopia

Background and Rationale

In Ethiopia, the Amhara Region has both the highest known prevalence of active trachoma in children one to nine years, as well as TT in adults ages 15 years and older. While there have been successes in the S and A components of the SAFE strategy, including high, validated drug coverage, the region has not yet reached the UIGs. MDA alone is likely not enough to eliminate trachoma in areas which are most affected by the disease and facial cleanliness and environmental improvement must be included in elimination strategies. Understanding the barriers to F and E uptake is crucial in meeting the UIGs in Amhara.

Aim

This study aims to understand the behavioral factors that promote and inhibit good hygiene practices, such as facial cleanliness and latrine usage, across the Amhara Region in order to develop new trachoma health education messages and tools.

Study Objectives

1. To assess important behavioral determinants influencing face washing of children
2. To assess important behavioral determinants influencing building and repair of latrines and latrine usage
3. To explore existing opportunities for behavior change and communications (BCC) in the Amhara Region and determine appropriate messages and tools for teaching and for both the F and E components of SAFE

Methods

For Objectives One and Two, a cluster/non-doer survey will be used to identify the main determinants that influence the face washing behaviors of mothers of children

Sample Size

The proposed sample size for the doer/non-doer survey can be seen in Figure 1 below.

Figure 1. Sampling Matrix for Doer/Non -Doer Surveys

Behavior	Priority Group	No. of Study Sites for each survey	Minimum No. of Participants
Face washing	Mothers of children ages 1-6 years		

Progress on WASH for Trachoma Elimination

Presented by Yael Velleman, Senior Policy Analyst (Health and Hygiene), WaterAid, speaking on behalf of the ICTC WASH Working Group and WHO

Background

In the past few years, there has been increased recognition of the crucial role of WASH in the effective control of blinding trachoma to achieve the 2020 elimination target. This recognition is leading to increased collaboration with WASH stakeholders, as well as moving beyond coordination to improved practice. This presentation outlines some of the key activities currently under way in order to receive feedback from meeting participants and engage them in forthcoming processes.

Strategy and Guidance

WHO

At the 2014 WHO Alliance for the Global Elimination of Blinding Trachoma by the year 2020 meeting, the need for increased internal coordination within WHO on WASH and trachoma was noted as a key consideration. WHO has since set up an internal coordination group bringing together the NTDs department and the Water Sanitation and Health team. This group has initiated a process of defining an organizational strategy for WASH and NTDs, with trachoma as one of the focus diseases. The draft strategy will be finalized in the next few

Over 2013-2014, there have been several intensive formative research studies focused on identification of barriers and norms around F and E interventions in Northern Kenya and Uganda.

- Research to inform the development of behavior change interventions for the F and E of the SAFE strategy in Turkana and Marsabit, Kenya – November 2013 – London School of Hygiene and Tropical Medicine.
- Understanding Individual and Contextual Factors for Development of a Behavior Change Communication Campaign for Trachoma Prevention In Busoga and Karamoja Regions, Uganda – April 2014 – Johns Hopkins University Center for Communications Programs.

The Fred Hollows Foundation, in collaboration with other ICTC members, is undertaking work to address specific knowledge gaps surrounding two areas: the behavioral drivers of facial cleanliness and other hygiene practices to inform better behavior change programming (through funding formative research work in Ethiopia) and the cost of implementing F and E for trachoma elimination in national programs at different

International Coalition for Trachoma Control Update

Presented by Dr. K.H. Martin Kollmann, Chair of the ICTC

Background

The trachoma community is making great progress as demonstrated by the second progress report on the London Declaration on NTDs, which tracks progress against landmark commitments made in 2012. The 2014 report, produced by United against NTDs, indicates that trachoma is on track against all indicators, a great encouragement to those involved in the fight against trachoma. Further the WHO Weekly Epidemiological Report demonstrates that trichiasis surgeries and antibiotic distributions have increased in significantly over the last decade even though the challenge of implementing F and E components remains significant.

This success is building on:

- A strong global partnership (WHO GET2020 Alliance)
- Local ownership and leadership (national programs)
- An agreed public health approach (SAFE strategy)
- A drug donation program (International Trachoma Initiative (ITI))
- The support of the ICTC
- Donor interest and coordination

Progress in 2014

Working Groups

ICTC's seven working groups launched in early 2014 are the primary channel for member engagement and member ci EMC310050F(a)12(cis)8(s)-4toon the on e(k c)6(nal)i2(cnt.)] TJ ET BT 1 0 0 2 392524 336.41 T

Grasping the Scale of the Problem

The GTMP, which aims to survey every suspected endemic district in the world, made exceptional progress in 2014. The project met its original target well ahead of schedule in September 2014. By the end of 2014, GTMP had mapped 94 percent of the targeted districts and examined over 1.8 million people in 21 countries.

Funding Streams

Diverse, generous funders support SAFE strategy implementation including USAID, DFID, Lions Clubs International Foundation, The Queen Elizabeth Diamond Jubilee Trust, Pfizer, Conrad N. Hilton Foundation, and others. The following grants are managed directly by ICTC as a consortium, also known as 'partner initiatives': The Queen Elizabeth Diamond Jubilee Trust Trachoma Initiative and DFID SAFE. A number of ICTC members also work together on the GTMP supported by DFID, through a grant to Sightsavers and USAID.

ICTC Partnership Initiatives

In 2014, ICTC members together with funding partners launched two large scale trachoma elimination initiatives: The Queen Elizabeth Diamond Jubilee Trust Trachoma Initiative in Africa, which is working in Uganda, Kenya, Malawi, Mozambique, Nigeria, and Tanzania, and DFID's Trachoma SAFE Implementation Program currently working in Chad, Ethiopia, Tanzania, and Zambia.

Other Initiatives

- Trust Trachoma Initiative in the Pacific has invested in ICTC membership. The Fred Hollows Foundation, who has reinvested funds in working with The Queen Elizabeth Diamond Jubilee Trust, local government departments and NGOs, and nongovernmental organization partners to identify the programming mix required to eliminate blinding trachoma in four Pacific Island nations.
- Eastern Mediterranean Region Alliance for Trachoma Control also has sourced some funding. ICTC has been working with the Alliance through a memorandum of understanding in supporting capacity building workshops, sharing technical resources and engaging in practical field experiences. ICTC and Alliance members have also supported GTMP mapping in the region and are coordinating a funding proposal for full SAFE implementation in Egypt, Pakistan, and Sudan.
- ICTC member HKI was appointed the prime partner in a five-year, $35 million Cooperative Agreement Grant on Morbidity Management and Disability Prevention for blinding trachoma and Lymphatic Filariasis activities. The other partners are RTI International, Filariasis Morbidity Project, and the Kilimanjaro Center for Community Ophthalmology.

Key Challenges

WASH

ICTC is working closely with members and other stakeholders, including WHO, on articulating WASH strategies and principles and defining practical indicators to address F and E.

Scale up and capacity-building

Scale up requires coordinated capacity-building around cost saving, coordination, and translation of global policy to local practice and program management.

More Funding Needed

In 2011, our ICTC global roadmap 2020 INSight estimated that 437-748 million USD would be needed to reach elimination. Significant scale up and finalizing the fight against trachoma brings new challenges affecting cost.

Upcoming Highlights

- Post-2015 strategy planning: ICTC is undertaking a strategy planning consultation to develop its post-2015 strategy in response to a rapidly changing environment and challenges ahead.
- Advocacy resources for 2015: To advocate for the support needed to meet the 2020 elimination deadline, ICTC is developing a number of key tools, including a gap analysis to highlight unfunded countries and areas (in collaboration with WHO), a 'Global SAFE Implementation Calculator' to help identifying the cost of elimination in different environments, and a progress report on the 2020 Insight: the end in sight global roadmap.
- New Vice Chair elections: ICTC will welcome a new Vice Chair in April 2015.

Facial Cleanliness and Environmental Improvement (F and E) Indicators: Showing Impact through Purpose

Presented by Angelia Sanders, Associate Director of the Trachoma Control Program, The Carter Center

Background

iv. Facility is shared among no more than five families or 30 persons, whichever is fewer, and if the users know each other

3. *100 percent of latrine coverage in schools*
    a. Latrine coverage is defined as:
        i. At least one toilet per 25 girls and at least one toilet for female staff
        ii. A minimum of one toilet plus one urinal (or 50 centimeters of urinal wall) per 50 boys and at least one toilet for male staff
        iii. Is equipped with hand/face washing stations inside or immediately outside the sanitation facility

4. *100 percent latrine use*
    a. Latrine use is considered met when at least *three of five* of the below signs (or other culturally appropriate signs) are present:
        i. Path to latrine worn
        ii. Presence of fresh feces in latrine
        iii. Water in/next to latrine used for anal cleansing and/or hand washing
        iv. HH verbally confirms use
        v. No feces observed in compound and within vicinity of compound (Open Defecation Free)

5. *100 percent of HHs have access to protected water source*
    a. Protected water sources include:
        i. Shallow well
        ii. Deep well
        iii. Borehole pump
        iv. Rain catchment
        v. Piped water

These indicators were categorized based on the need to answer six questions (see Table 1). The answers will provide programs with a wealth of information, such as whether children's' faces are clean; households have and are using latrines; children have access to latrines while in school; and if households have access to water. Clean faces, access to and use of latrines both at home and in school, and access to adequate all water are potential contributors to reduce risk of infection. In the case of children, having access to latrines in school further reinforces latrine use behavior change and health education messaging. Additionally, due to the fact that many of these indicators overlap with current indicators used by the WASH sector, it should be easier for the different sectors to work together, streamline activities and share data.

F and E Activities

In order for programs to successfully reach their indicators, activities must be conducted and documented. There are many activities that ministries of both health and water can conduct in order to reach the indicator goals; however, not all of them are easy to document on a regular basis. Recognizing the difficultly in reporting at the village level, four primary activities (see Table 2) are recommended as actions to be taken and documented in monthly community health workers' (CHWs) reports. These in 1 72.024 217.91.04 Tf 1 0 0 1 97.584 4

Table 1 Proposed Indicators

What (indicator/goal)	100% Children 19	

Table 2 Activities in the Field

What (indicator/goal) Who (unit of measurement)	HE Sessions conducted	New latrines provided	New Water sources Constructed	Water sources repaired

New WHO Recommendations for Trachoma Surveillance

Presented by Dr. Anthony Solomon, Medical Officer, Neglected Tropical Diseases, World Health Organization

Background

The new WHO Trachoma Elimination Program Standar

Zithromax® Global Supply

Presented by Julie Jenson, Director Corporate Responsibility/Brand Supply Lead Zithromax®, Pfizer Inc.

Background

Since the London Declaration launched in 2012, because of commitments made by donors and ministries of health, Pfizer has increased its donation of the antibiotic Zithromax® from 50 million treatments in 2014 to a projected 80 million treatments in 2015. Sixteen years ago, Pfizer made the commitment to provide Zithromax®

University of California San Francisco Carter Center Research Partnership

Presented by Nicole Stoller, Program Manager, Francis I. Proctor Foundation

Background

UnWIo

treatment of pre-school children was performed, and from 2.8 percent to 7.0 percent where targeted treatment of clinically active cases and their households was completed.

Please note: all PCR results are expressed as a percentage of positive pools, and are preliminary pending the completion of individual sample processing.

MORDOR: Mortality Reduction After Oral Azithromycin

An estimated 7.7 million pre-school aged children die each year, the majority from infectious diseases.[12] Recently, a 2011 study that the group conducted demonstrated that mass azithromycin distributions for trachoma may have the unintended benefit of reducing childhood mortality.[13][14] This surprising result was observed in an area of Ethiopia with highly prevalent trachoma. Another trial is necessary to determine whether a similar effect will be seen in areas not eligible for trachoma treatments. If mass azithromycin can be proven to reduce childhood mortality, then widespread implementation in developing countries could be

East Africa: Kilosa district, Tanzania (Tanzania National Institute for Medic

MDA for Trachoma Control: Looking Back, Moving Forward

Violeta Jimenez, Consultant, Emory University
Aisha Stewart, Associate Director, Trachoma Control Program, The Carter Center
Dr. Scott Nash, Epidemiologist, Trachoma Control Program, The Carter Center

Looking back, a considerable body of evidence has been generated on the use of Zithromax® for trachoma control. In highly endemic areas, such as in the Amhara region of Ethiopia, the prevalence of TF has declined after intensive control efforts with Zithromax®. However despite tens of millions of doses of

Discussion Points

Given the results of the many community randomized trials, and what we know to date about MDA for trachoma control, it is important to take a step back and have a conversation among the wider trachoma community to help set priorities and goals. Some important questions to be discussed by the global community include:

- At what point do we have enough evidence to take action (at a programmatic level) and who decides?
- What is feasible to implement at scale?
- What resources are required?
- How long can an approach be sustained?
- Can study results be applied in different settings?
-

The Impact of Trichiasis on Quality of Life and Poverty

Presented by Matthew Burton, Senior Lecturer and Wellcome Trust Senior Research Fellow, London School of Hygiene and Tropical Medicine

Background

Trachoma is considered a disease of poverty that significantly affects individuals' quality of life. Although there are many epidemiological studies linking trachoma to factors normally associated with poverty, formal quantitative data linking trachoma to household economic poverty in endemic communities is relatively limited. Likewise, little formal quantitative data support the conclusion that TT affects individuals' quality of life. This dual study aimed to show the relationship between trichiasis and poverty and also trichiasis and quality of life.[15,16]

Aim

To determine the relationship between trichiasis and economic poverty and trichiasis and the quality of life.

Methods and Results

Trichiasis and Poverty

Two hundred people with TT were recruited through community-based screening. Cases were individually matched by age and gender to 200 controls without trichiasis, selected randomly from the same village as the case. A broad set of asset-based indicators was measured and household economic poverty determined

0 a1 0 tiveBul-e 0

selected randomly from the sub-villages of every fifth trichiasis case were matched by age (within five years) and sex.

Vision and health related quality of life was measured using the WHO/PBD-VF-32 and WHOQOL-BREF tools respectively. Data on the impact of TT on social relations, marriage, and housekeeping were collected through semi-structured questions. The comparison of the quality of life scores between cases and controls were made using linear regression adjusted for age, sex, and socioeconomic status.

The TT cases had substantially lower vision and health related quality of life in all domains than controls (P<0.0001). Trichiasis cases also had significantly lower overall quality of life score (Mean score: 34.5 v 64.6, p<0.0001) and overall health satisfaction (38.2 v 71.7, p<0.0001) than the controls. In trichiasis cases, longer trichiasis duration, central corneal opacity, poor vision, and contrast sensitivity scores were independently associated with low vision related quality of life in all subscales. Not having partner (p<0.0001), daily laboring (p<0.0001), unemployment (p=0.0010), poor vision (p 0.0076), poor contrast sensitivity (p 0.002), and presence of other health problems (p<0.0001) were independently associated with poor overall health related quality of life score in trichiasis cases. Due to their trichiasis: 596 (59%) feel ashamed of or embarrassed; 913 (91.3%) worry that they will lose their remaining eye sight; 701 (70%) have been troubled in their marriage and ignored by their marriage partners; and 681 (68%

Table 1. Surgical quality assessment results, Kayes region, Mali, 2014.

District	Date Surveyed	# Patients Selected	# Patients Visited

13. Programs should ensure funding to carry out reverse drug logistics or transportation of Zithromax® within a country to ensure all stock is used before expiration and to ensure all countries maximize the use of the limited supply.

14. Countries conducting TT surgeries should carry out validation studies to verify reporting.

15.

Trachoma: The Disease

Trachoma, the world's leading cause of preventable blindness, can be found in over 50 countries. More than 232million people are at

Tuesday, March 3	
7:30	*~Depart the Sheraton Hotel for The Carter Center~*
8:00– 8:30	[redacted]

Wednesday, March 4		
7:30	*~Depart the Sheraton Hotel for The Carter Center~*	
8:00– 8:30	Breakfast	
8:30– 8:45	Chairperson Announcements	Dr. Emmanuel Miri Country Representative The Carter Center Nigeria
8:45– 9:30	Ethiopia S Presentation & Discussion	Mr. Oumer Shafi NTD Team Leader Federal Ministry of Health Ethiopia
9:30– 10:00	The impact of trichiasis on quality of life a poverty	

Ethiopia
Mr. Ali Gebeyehu (ARHB)
Mr. Oumer Shafi Abdurahman (FMOH)
Dr. Zerihun Tadesse (The Carter Center)
Mr. Belay Bayissasse (The Carter Center)
Mr. Mohammed Aderajew (The Carter Center)
Mr. Eshetu Sata Shanka (The Carter Center)
Mr. Mulat Zerihun Lemu (The Carter Center)
Ms. Tara Brant (The Carter Center)

Mali
Prof. Lamine Traoré (MOH)
Mr. Sadi Moussa (The Carter Center)
Mr. Yaya Kamissoko (The Carter Center)

Niger
Dr. Kadri Boubacam [(T 1 0a-6.98 1 0 0 1 89.447 Tm [(Dr)-4(.)] TJ 159.7T 1 0 0 1 89.4(o)10(r)é)4((M)5(OH